

Rapid #: -18353730

CROSS REF ID: **11290117100005153**

LENDER: **UNL :: Main Library**

BORROWER: **CEF :: Main Library**

TYPE: Article CC:CCG

JOURNAL TITLE: Nature reviews. Earth & environment

USER JOURNAL TITLE: Nature Reviews Earth & Environment

ARTICLE TITLE: A low-to-no snow future and its impacts on water resources in the western United States

ARTICLE AUTHOR: Siirila-Woodburn, Erica R;Rhoades, Alan M;Hatchett

VOLUME: 2

ISSUE: 11

MONTH:

YEAR: 2021

PAGES: 800-819

ISSN: 2662-138X

OCLC #:

Processed by RapidX: 12/1/2021 12:10:52 PM

This material may be protected by copyright law (Title 17 U.S. Code)



A low-to-no snow future and its impacts on water resources in the western United States

Erica R. Siirila-Woodburn ^{1,9}✉, Alan M. Rhoades ^{1,9}, Benjamin J. Hatchett², Laurie S. Huning ^{3,4}, Julia Szinai ^{1,5}, Christina Tague ⁶, Peter S. Nico ¹, Daniel R. Feldman ¹, Andrew D. Jones ^{1,5}, William D. Collins ^{1,7} and Laurna Kaatz⁸

Abstract | Anthropogenic climate change is decreasing seasonal snowpacks globally, with potentially catastrophic consequences on water resources, given the long-held reliance on snowpack in water management. In this Review, we examine the changes and trickle-down impacts of snow loss in the western United States (WUS). Across the WUS, snow water equivalent declines of ~25% are expected by 2050, with losses comparable with contemporary historical trends. There is less consensus on the time horizon of snow disappearance, but model projections combined with a new low-to-no snow definition suggest ~35–60 years before low-to-no snow becomes persistent if greenhouse gas emissions continue unabated. Diminished and more ephemeral snowpacks that melt earlier will alter groundwater and streamflow dynamics. The direction of these changes are difficult to constrain given competing factors such as higher evapotranspiration, altered vegetation composition and changes in wildfire behaviour in a warmer world. These changes undermine conventional WUS water management practices, but through proactive implementation of soft and hard adaptation strategies, there is potential to build resilience to extreme, episodic and, eventually, persistent low-to-no snow conditions. Federal investments offer a timely opportunity to address these vulnerabilities, but they require a concerted portfolio of activities that cross historically siloed physical and disciplinary boundaries.

Mountains are known as the water towers of the world, capturing, storing and releasing water for downstream use^{1,2}. In the western United States (WUS), as in many global regions², this natural service largely occurs through seasonal mountain snowpacks (FIG. 1a), storing approximately 200 km³ or 162 million acre-feet (MAF) of water annually^{3,4}, as quantified through snow water equivalent (SWE). In places such as California, for example, average 1 April snowpack water storage (21 km³ or 17 MAF) nearly doubles the amount of surface water reservoir storage (29 km³ or 23.5 MAF)⁵. The springtime and summertime melt of these seasonal snowpacks is fundamental to water infrastructure and operations, supplying water during times of precipitation scarcity and when agricultural, ecological and municipal water demands are high⁶. Over the last century, observations and models reveal that anthropogenic climate change has substantially reshaped WUS water resources, including a declining snowpack^{7–9,10–17}. For instance, observed 1 April snowpack decline since the mid-twentieth century ranged between 15 and 30% or 25 and 50 km³ (REF.¹⁵).

Future mountain snowpacks are further projected to decline, and even disappear, but at unknown rates⁴. While the complete loss of snow is the worst-case scenario, a plausible situation informed by estimates of historical low-snow conditions (FIG. 1b) would be a WUS-wide reduction in SWE and seasonal snow, and a shift from rare or short-term to more persistent low-to-no snow occurrences.

The potential for persistent low-to-no snow to disrupt the WUS water system is substantial, potentially even catastrophic. Water storage and conveyance infrastructure was designed and is now managed using spring snowmelt as a central criterion for operations^{5,18–22}. These water management decisions are predicated on the assumption of a stationary climate^{19,23}, which is an unintended, yet, critical, oversight. Moreover, water rights allocations for many major WUS rivers were made in the late nineteenth and early twentieth century at the end of the Little Ice Age (AD 1400–1900)^{24–28}, a period that is among the wettest in the past 4,000 years^{29–32}. Given the high confidence that continued warming will result in

✉e-mail:
ERwoodburn@lbl.gov
<https://doi.org/10.1038/s43017-021-00219-y>

Key points

- Mountain snowpacks in the western United States (WUS) have historically acted as large, natural reservoirs of water; yet, they are now harbingers of a changing climate through their signalling of a low-to-no snow future.
- Models projecting the time horizon of low-to-no snow in the WUS lack spatiotemporal consensus due to differences in definitions, metrics, methods and regionally specific analyses.
- Low-to-no snow will impose a series of cascading hydrologic changes to the water–energy balance, including vegetation processes, surface and subsurface water storage and, ultimately, streamflow that directly impacts water management.
- A re-evaluation of long-standing hydroclimatic stationarity assumptions in WUS water management is urgently needed, given the impending trickle-down impacts of a low-to-no snow future.
- Observational and modelling advances are needed to better understand the implications of a low-to-no snow future on water resources and to evaluate the trade-offs among a wide array of potential adaptation strategies that can address both water supply availability and water demands.
- Co-production of knowledge between scientists and water managers can help to ensure that scientific advances provide actionable insight and support adaptation decision-making processes that unfold in the context of significant uncertainties about future conditions.

decreased river flows in the WUS, of which snow-loss is a major contributor, warming will challenge the ability to meet water demands under allocations that did not include climate change risk management¹⁷.

Societies established on freshwater availability and seasonal storage capacity in mountainous snowpack could, therefore, face multibillion-dollar implications, with costs dependent on the time horizon of a low-to-no snow transition⁴. For example, assuming a WUS-wide loss of 66.6 km³ or 54 MAF of annual snow storage valued at \$200 per acre-foot, with low-to-no snow conditions emerging in the next 50–100 years, WUS snow loss is estimated to reach a cumulative cost of US\$120–850 billion⁴. Of course, economic impact assessments of snow loss are complicated by water values that depend on time, location and supply abundance, as well as the nonlinear changes in the value of water during severe drought. Yet, despite these complications, such estimates demonstrate the dire situation of a low-to-no snow future with little to no action.

Uncertainties in the time horizon of snow loss also have practical implications and challenges for the diverse set of water users, including for the agricultural

and municipal sectors^{33,34}. Future snowpack losses will be detrimental to long-term water stores typically reserved for emergencies^{35,36} and become increasingly difficult to manage as transitions between low-to-no snow periods move from extreme (single year), to episodic (multi-year), to persistent (decades) conditions. Moreover, the disappearance of snow in the WUS has important hydrologic ramifications on both natural and managed systems. Changes in the seasonal snow cycle influences the timing and magnitude of ground-water recharge, vegetation dynamics and stream discharge, which then directly impacts water availability. These processes occur simultaneously and demonstrate nonlinear and heterogeneous behaviour^{37–40}.

Thus, there is a real need to understand the time horizons, spatial extents and magnitudes of snowpack changes to inform what can be gained through climate change mitigation. This information needs to be provided at decision-relevant scales and conducted with use-inspired science that is informed by a diverse stakeholder group^{41,42}. Ultimately, water policy and governance needs to be constrained by the spatiotemporal limits of the water cycle to avoid the prior pitfalls of assuming climate stationarity. This approach of being proactive rather than reactive to a low-to-no snow future will safeguard already limited natural and financial resources, and aid in building resilience to a low-to-no snow future.

In this Review, we provide a call to action and forewarn the dire implications of a low-to-no snow future, given its central role in mountainous watershed behaviour, ecosystem function and, ultimately, downstream water availability. We begin by synthesizing observational evidence of snowpack disappearance in the WUS, followed by the propagating impacts of those changes on the hydrologic cycle. We follow with discussion of adaptation strategies necessary to reduce the economic and ecosystem impacts of low-to-no snow. We end with recommendations for future research, including the need for concerted efforts to cross historically siloed physical and disciplinary boundaries, as well as the scientific practitioner divide. While other hydro-meteorological variables have important bearing on WUS water resources, emphasis is placed on snowpack changes, owing to the historic reliance on snowpack by water resources management as a seasonal natural reservoir, and due to a strong, inverse relationship with temperature¹⁷ compared with precipitation^{43–45}, which lacks consensus on whether future conditions in the WUS will be wetter, drier or remain the same.

A declining mountain snowpack

Prior to assessing future projections of WUS snowpacks in the face of anthropogenic climate change, it is crucial to provide context on the historical and contemporary changes of snowpack and the observational networks used to measure them.

Observational networks and their limitations.

Observational networks of WUS snowpack are some of the most extensive in the world, including centennial-length snow courses, decadal-length manual and

Author addresses

¹Earth and Environmental Sciences Area, Lawrence Berkeley National Laboratory, Berkeley, CA, USA.

²Western Regional Climate Center (WRCC), Reno, NV, USA.

³Department of Civil Engineering and Construction Engineering Management, California State University, Long Beach, Long Beach, CA, USA.

⁴Department of Civil and Environmental Engineering, University of California, Irvine, Irvine, CA, USA.

⁵Energy and Resources Group, University of California, Berkeley, Berkeley, CA, USA.

⁶Bren School of Environmental Science & Management, University of California, Santa Barbara, Santa Barbara, CA, USA.

⁷Department of Earth and Planetary Science, University of California, Berkeley, Berkeley, CA, USA.

⁸Denver Water, Denver, CO, USA.

⁹These authors contributed equally: Erica R. Siirila-Woodburn, Alan M. Rhoades.




Fig. 1 | Spatiotemporal variability in western United States seasonal snowpack. a | Median peak snow water equivalent (SWE) for seasonal snowpacks of the western United States over 1982–2016 (REF.²⁵⁵). California and Upper Colorado hydrologic units, as defined by the United States Geological Survey, are distinguished by green and blue contours, respectively. **b** | As in panel **a**, but the minimum peak SWE for seasonal snowpacks. **c** | The temporal evolution of mean and historic high-snow and low-snow water years (WY) for the California hydrologic unit (left), and the cumulative distribution function of peak SWE water storage volume (right). Vertical bars in the right panel indicate peak SWE and median SWE for the time series illustrated in the left panel. **d** | As in panel **c**, but for the Upper Colorado hydrologic unit. Interannual differences in SWE conditions can proximally serve as an indicator of future low-to-no snow conditions.

automated in situ measurements, and multi-year airborne and remotely sensed products.

An important observation date for WUS snow observations is 1 April. While somewhat arbitrary, this date results from monthly manual snow surveys established

early in the twentieth century^{46,47}, and is codified into water management as an indicator of warm season streamflow³⁵. However, depending on location and hydroclimatic variability⁴⁸, 1 April SWE does not necessarily represent the total amount of water stored as

snow in a given water year⁴⁹ (FIG. 1c,d), nor can it capture the warming-induced earlier shift in the peak timing of SWE^{50,51}. Peak SWE can better quantify the total amount of meltwater available and identify when the melt season might begin, but it cannot fully characterize the temporal evolution and spatial variation of annual water storage and release^{12,29,52,53}, especially in lower elevation, ephemeral snowpacks that have multiple peaks in the water year life cycle of SWE⁵⁴. In the Rocky Mountains, 1 April SWE underestimates peak SWE by 12%⁵⁵. Both variables are, thus, often needed to assess whether snowpack changes represent an absolute loss or a temporal shift.

Other metrics are also used to characterize SWE. SWE centroid accounts for coarser temporal data (monthly) and snowpacks with multiple SWE values⁵⁰, but it inherently provides a more constrained estimate of meltwater totals and can artificially influence other management-relevant metrics, such as the snowmelt season start date, snowmelt season length and/or spring snowmelt rates¹⁸. SWE centroid and peak SWE also fail to account for SWE changes at daily-to-subseasonal scales⁵⁶. ‘Snow drought’ refers to a historically low snowpack in a particular time in the water year, or over multiple water years, created by either anomalously low precipitation (dry snow drought) and/or anomalously high temperatures during precipitation events or driving midwinter melt (warm snow drought)^{29,57,58}. The concept has gained momentum and enables a better understanding of the drivers of low-to-no snow conditions and the identification of when snow conditions begin to diverge from normality.

Interannual variability that drives differences between peak SWE and 1 April SWE, as well as changes in snow accumulation and persistence as a result of anthropogenic climate change, highlight the importance of observational networks in providing real-time monitoring of current snowpack conditions. Yet, the utility of these observational networks is limited by instrument measurement biases and spatiotemporal gaps^{59–61}. A spatially incomplete in situ network, especially at high-elevation headwater regions, creates challenges as warming continues to drive the freezing line upslope, which inhibits the accumulation and persistence of seasonal snowpack, particularly at mid-elevations^{14,62,63}. Loss of snowpack can further reduce the ability of in situ networks to provide skilful drought prediction, given a strong relationship between snowmelt and runoff²².

To address challenges associated with monitoring gaps and extreme events, various statistical and model-based interpolation methods have been developed, so as to provide spatially distributed information. However, assumptions made in interpolation methods produce 40–66% differences in estimates of WUS-wide peak SWE, limiting direct applications to water resource management⁶⁴. Blending traditional approaches, such as snow sensors and expanded snow course surveys⁴⁷, with novel techniques like airborne lidar⁶⁵, remotely sensed observations^{53,66,67} and citizen science⁶⁸, can offset some of these limitations. Nevertheless, heterogeneity in mountainous environments limit the ability to obtain perfect snowpack observations, necessitating the use of models⁶¹.

Estimates of change. Despite observational challenges, estimates of contemporary changes have been widely documented, revealing significant reductions in WUS mountain snowpacks and their persistence (FIG. 1c,d). Indeed, observed reductions in both peak SWE and 1 April SWE provide two lines of evidence for a declining snowpack^{15,69} (Supplementary Table 1). Peak SWE timing, for instance, has shifted 8 days earlier per degree of warming⁵¹, while peak SWE losses of -2.8 mm per year across 25% of stations have been estimated⁶⁹. Moreover, 1 April SWE depths have decreased by 21% across 90% of reporting stations over 1955–2016, representing a snow loss equivalent to the storage capacity of Lake Mead, the largest reservoir in the WUS¹⁵. Below-normal conditions have further increased in likelihood in sequential years^{1,51,57,70,71}.

While WUS mountains are dependent on extreme precipitation and snowfall events to build snowpacks, they are also susceptible to changes in precipitation characteristics, contributing to some of the observed changes alongside warming. Atmospheric rivers, for example, which are commonly associated with snowpack accumulation, are warming, intensifying and producing more rainfall and thus, rain-on-snow events^{35,36,72–74}. Since the 1980s, rates of minimum temperature warming in WUS mountains ranges have increased from 0.17 to 0.7 °C per decade, with greater warming (0.8–1.2 °C per decade) across more locations on wet days^{72,75}. Warming temperatures push the snowline upslope, nonlinearly altering the amount of water that can be stored as snow^{63,76}. Warming also causes snowpacks to ripen earlier in the season, increasing snowpack susceptibility to rain-on-snow events⁷⁷. Deposition of light-absorbing particles — including dust and black carbon — further lower snow albedo, increase solar energy absorption and accelerate snow depletion by 31–51 days, so as to result in earlier peak runoff timing^{60,78–80}. Increasing wildfire activity⁸¹ and land-use change⁷⁹ are enhancing the deposition of light-absorbing particles on WUS snowpacks, though regional impacts of these particles on the snowpack energy balance remain uncertain^{82,83}.

Given the strong reliance on an increasingly volatile source of water, resource managers in the last decade have proactively turned to innovative strategies to retain higher reservoir levels via advances in hydrometeorological forecasting⁸⁴. Low snowpack and, more broadly, drought have also motivated landmark policy changes, such as the Sustainable Groundwater Management Act in California⁸⁵.

A low-to-no snow future

With observed contemporary changes in the WUS snowpack and projected continuation of anthropogenic climate change, there is high-to-very high confidence that WUS-wide snowpack will continue to decline and peak earlier in the water year¹⁷. However, for various reasons, little consensus exists regarding when snowpack disappearance becomes particularly detrimental, when the time-to-emergence of low-to-no snow might arise and how the spatial extent of a low-to-no snow future will materialize, as will now be discussed.




Fig. 2 | **A multiscale perspective of the mountainous hydrologic cycle.** Earth system spatial scales (continental, mountain range and watershed) as they pertain to western United States water and their influences from physical processes (left) and management strategies (right), which also vary by scale. The dashed black box in the continental scale schematic is representative of a typical Earth System Model gridcell (~100 km), sometimes equivalent to an entire mountain range, encapsulating substantial land and subsurface heterogeneity. These cross-scale and cross-discipline considerations are required for a holistic assessment of water management in the western United States.

Challenges of producing snow projections. Snowpack is an emergent property of cross-scale interactions involving temperature, precipitation, radiation, topography and land-surface characteristics, all of which span spatial scales of the globe to hillslopes and temporal scales of centuries to seconds⁸⁶ (FIG. 2). For decades, a lack of cross-scale theory, as well as siloed sciences, sparse observational data and computational limitations, constrained advances in modelling mountain environments with high fidelity. Although Earth System Model (ESM) capabilities have dramatically evolved in the last century⁸⁷, particularly their strengths as large-scale, centennial-length projection tools, ESMs lack some key process representations needed to accurately simulate mountainous environments, particularly at decision-relevant scales (BOX 1). Indeed, a fully coupled bedrock-through-atmosphere system remains a scientific grand challenge, given limitations in resolving spatial and temporal heterogeneity of atmospheric and land-surface hydrological processes (ranging from tens to hundreds of metres), the substantial disparity in model scales between scientific communities (sub-kilometre to hundreds of kilometres) and the uncertainties around the physical coupling, or lack thereof, between models and/or model components (FIG. 2).

Recent breakthroughs in supercomputing (for example, Exascale and GPU computing)^{88–90}, benchmarking

of models to high-spatiotemporal-resolution observations (including remotely sensed and readily deployable in situ observational networks)^{91–94} and the growing field of artificial intelligence^{95,96} are accelerating ESM development and their ability to represent the mountainous hydrologic cycle. These breakthroughs combined with the use of variable and/or adaptive mesh refinements to aid in scale-awareness across resolutions^{97–103}, mountain-focused parameterizations^{104,105}, more flexible and modular codes^{106,107}, determination of when and where two-way coupling is needed between models and submodels¹⁰⁸, and utilization of machine learning to better constrain parameter uncertainty or lack of cross-scale theory⁹⁵ will inevitably reduce the wide range of estimates in future projections of mountain snowpack.

The use of a hierarchy of models, namely, intermediate-scale (single to tens of kilometres) integrated process models (IPMs), can further inform assumptions made in ESMs and their ability to project future snow loss. IPMs are computationally less expensive, can more realistically represent important model lower boundary features and can be more easily benchmarked with and/or periodically updated by observations^{109,110}. Most importantly, they can be used to interrogate the process assumptions made in ESMs by systematically removing process fidelity in the IPM for those found in ESMs. However, considerations

Box 1 | Mountainous hydrologic cycle process uncertainty in Earth System Models

The mountainous hydrologic cycle provides a unique test for Earth System Models. The mosaic of cross-scale interactions and emergent behaviour stress the validity of model assumptions, particularly subgrid parameterizations. Indeed, process representation in atmospheric, land-surface and subsurface systems — needed to characterize the partitioning, stores and fluxes of the mountainous hydrologic cycle — remains a grand challenge, specifically under impending low-to-no snow conditions. Earth System Model limitations include:

- Simplified microphysical and macrophysical representations of dominant hydrometeor processes that can bias the magnitude and elevational gradients in mountain precipitation^{256,257}, and often do not account for wind redistribution and sublimation during or after precipitation^{111,258}.
- Parameterization omission of terrain influences on shortwave and longwave radiation, which impacts the spatial heterogeneity and lifetime of snowpack^{259–261}.
- Assumptions in parameterization choice (such as aerodynamic roughness length) can decouple the atmosphere–land interface, particularly during the winter season, resulting in a rapid surface cooling effect that leads to large biases in surface temperature^{257,262}.
- Prescription of snowpack density and snow cover fraction by empirical functions based on discrete temperature ranges applied uniformly across mountain ranges, ignoring the influences of relative humidity²⁶³ and wind redistribution²⁵⁸, impacting surface albedo, water storage and energy fluxes.
- Under-representation of the multiple interacting gradients needed to resolve vegetation dynamics (for example, topographic, biogeographic, disturbance), impacting vegetation processes during cold season precipitation (such as canopy interception) and growing season evapotranspiration¹⁰⁷.
- Neglected or crude representation of subsurface processes, including vadose zone dynamics, lateral flow and geological heterogeneity, influencing infiltration–runoff partitioning and groundwater recharge²⁶⁴.

should also be made for how to minimize bias propagation between coupled models and benchmarks that ensure that added value is actually provided with more costly, complex simulations. Lastly, systematic model intercomparison projects, such as the Snow Model Intercomparison Project (SnowMIP)¹¹¹, will be pivotal in providing directions for other important community and model development needs.

Projections of mountain snowpacks, therefore, become a computational balance between limiting the physical process representation at a particular scale (model resolution versus subgrid-scale parameterization); the length of the simulation (a single water year versus multiple decades); the assessment of climate internal variability and statistical robustness (single versus multiple ensemble members); and/or the number of emissions scenarios assessed. These constraints yield an extensive and diverse literature, with a wide range of descriptive metrics, projection methods, time periods and regional specificity (Supplementary Table 1).

Literature synthesis of WUS snow projections. As a result of these limitations, particularly the difficulty in comparing different metrics and widespread use of a single, high-emissions scenario, syntheses of WUS snowpack projections must be taken with caution. Nevertheless, bringing together 18 analyses^{3,7,21,104,105,112–124} of snowpack changes across the WUS and four major WUS mountain ranges reveals continued snow loss (FIG. 3; Supplementary Tables 1,2).

In particular, WUS-wide average SWE decline (and their 95% confidence intervals, both rounded to the

nearest 5%) is $\sim 25 \pm 5\%$ by 2050, $\sim 35 \pm 10\%$ by 2075 and $\sim 50 \pm 10\%$ by 2100 (FIG. 3a). The large spread in projected changes at mid-century to end of century highlights the lack of consensus on the time to emergence of low-to-no snow. The spread also highlights challenges about how to best characterize SWE loss and standardize SWE projection methods, such that scientific findings can be transformed into actionable information for water managers adapting to the rapid rate of snow-loss brought about by anthropogenic climate change^{41,42,125,126}. This actionable information is particularly pertinent for the maritime mountains of the WUS (the Cascades and the Sierra Nevada) (FIG. 3b,d), where $\sim 45\%$ losses are expected by 2050, compared with 20–30% for the continental mountains (the Rockies and the Wasatch/Uinta) (FIG. 3c,e). Additional projections and analyses are also needed in understudied mountains within, for example, the Great Basin and the Colorado Plateau.

Defining low-to-no snow. Declining SWE on a more frequent, persistent and widespread basis^{3,7,9,113}, albeit with regional variabilities (FIG. 3), is, thus, a hallmark of a projected shift to low-to-no snow in the WUS. Specifically defining low-to-no snow, however, is challenging owing to a multitude of considerations, including the metrics available to quantify snowpack decline.

In developing a quantitative definition of low-to-no snow, two critical points emerge. First, the definition should characterize conditions under which low-to-no snow becomes physically meaningful. Second, the snow conditions should represent deviations from ‘normal’ that result in challenges or failures in water management, recreational value and/or ecosystem services. The definition needs to be both broadly understood and regionally specific, while remaining applicable across WUS mountains and downstream water management paradigms. Precise definitions of low-to-no snow depend on the physical characteristics of snowpack (for example, volume of water stored in seasonal and ephemeral snow, peak timing of water storage, and runoff rate and timing) and the dependence of historic management paradigms.

From a physical perspective, low-to-no snow can be defined by its lowest bound (zero snow). Although not impossible, it is unlikely that a complete disappearance of snow in the WUS will occur before the end of the twenty-first century, even under a high-emissions scenario (FIG. 3), and deleterious impacts to resource management and ecosystems would likely occur before zero-snow conditions. Alternatively, a definition utilizing historic percentiles of peak SWE, akin to the US Drought Monitor¹²⁷, has been successfully applied to identify snow drought conditions^{9,12}. This approach provides a guide to defining low-to-no snow when combined with the strengths of the peak SWE metric. Within this percentile context, low snow is defined as conditions in which peak SWE falls between the 30th and 10th percentiles, and (effectively) no snow is defined as peak SWE conditions within the lowest 10th percentile.

Emergence of low-to-no snow. ESMs can be used to explore the time to emergence of low-to-no snow. They simulate the spatiotemporal scales necessary to assess





Fig. 3 | Ranges of projected twenty-first century snowpack loss. a | Projected snow water equivalent (SWE) loss across the mountainous western United States for the near future (2025–2049; yellow), mid-century (2050–2074; orange) and end of century (2075–2099; red), as derived from published literature^{3,7,21,104,105,112–124}. Percent SWE loss considers 1 April, peak SWE and/or seasonal SWE. Projection methods include Earth System Models (ESMs), bias-corrected statistically downscaled ESMs (BC-SD) and regional climate models (RCMs), incorporating analyses under RCP4.5 and RCP8.5, and, in some instances, older high-emission scenarios (SRES A2) (Supplementary Information). Box plots represent the standard minimum, maximum, upper and lower quartiles, and median projected SWE changes. **b** | As in panel **a**, but for the Cascades. **c** | As in panel **a**, but for the Rockies. **d** | As in panel **a**, but for the Sierra Nevada. **e** | As in panel **a**, but for the Wasatch/Uinta. The map at the top right illustrates the areas considered for the regional analyses, with ‘western United States’ encapsulating the entire domain. Heterogeneity in projected snowpack changes exists across mountain ranges and for different modelling approaches, but generally indicate agreement in decreases by the end of the century.



the timing, amount, phase and location of storms that deliver precipitation⁸⁷; capture alterations in the surface energy budget and the fluxes and stores of water; and avoid limitations associated with assumptions of stationarity¹⁹ through the use of emissions scenarios and land-use change. While acknowledging the outstanding limitations in ESM process representation in mountains (BOX 1), the uncertainty in individual

model projections¹²⁸, the value of large ensembles¹²⁹ and the varying plausibility of emission scenarios¹³⁰, exploration of the timeline to low-to-no-snow emergence is valuable, even within a single simulation as a proof of concept. Here, a 20-km-resolution simulation spanning 1950–2099 (REF.¹³¹) as part of the High Resolution Model Intercomparison Project (HighResMIP)¹³² is used for that purpose, adopting the

◀ Fig. 4 | **Snow disappearance based on a low-to-no snow definition.** **a** | Temporal evolution of the Upper Colorado hydrologic unit (see map on the right) categorized by peak snow water equivalent (SWE) percentiles. The white line indicates the 10-year running average of basin area (both snow and non-snow gridcells) considered low-to-no snow, that is, where an individual year's peak SWE \leq 30th historical peak SWE percentile. Data are from a single ensemble member of the 20-km-resolution MRI-AGRCM3-2-S Earth System Model¹³¹ under RCP8.5 (Supplementary Information). Vertical dashed and solid lines indicate the time-to-emergence of episodic and persistent low-to-no snow conditions, respectively, as defined by the onset of 5 and 10 sequential years with low-to-no snow conditions over \geq 50% of the basin. **b** | As in panel **a**, but for the Great Basin. **c** | As in panel **a**, but for the Pacific Northwest. **d** | As in panel **a**, but for California. Although illustrative of a single simulation, projections such as these demonstrate the utility of the 'low snow' and 'effectively no snow' definitions, as well as the time-to-emergence concepts.

\leq 30th percentile of historical peak SWE low-to-no snow definition (FIG. 4).

Through the middle and end of the twenty-first century, an increasing fraction of the WUS is impacted by SWE deficits relative to the historical period (FIG. 4; Supplementary Fig. 1). In particular, only 8–14% of years are classified as low-to-no snow over 1950–2000, compared with 78–94% over 2050–2099. In all regions, an abrupt transition occurs in the mid-to-late twenty-first century. For example, the onset of episodic low-to-no snow — the first time that five consecutive years of \geq 50% of the basin area experience low-to-no snow — occurs in the 2060s for most basins, but in California, emerges in the late 2040s. Persistent low-to-no snow — the first time that 10 consecutive years of \geq 50% of the basin area experience low-to-no snow — occurs as early as the late 2050s in California and as late as the end of the 2070s in the Upper Colorado (FIG. 4). Indeed, from 2070 to 2099, 80–97% of years meet the definition of persistent low-to-no snow. Projections indicate that situations can also arise where the shift from extreme to episodic conditions is rapid, such as in the Pacific Northwest, essentially skipping the episodic low-to-no snow regime¹³³. Therefore, if global emissions continued unabated, there is ~35–60 years before low-to-no becomes persistent across the WUS.

Propagating impacts to the hydrologic cycle

In a low-to-no snow climate, several key hydrologic processes controlling the spatiotemporal partitioning of water will be altered (FIG. 5), including precipitation stores (surface water, soil moisture and groundwater) and fluxes (evapotranspiration (ET), runoff and streamflow). Furthermore, deviations from 'normal' snow years can have direct and indirect impacts on hydrologic processes. These impacts are now discussed.

Evapotranspiration. Snowmelt is a significant source of water for vegetation in the WUS¹³⁴. Changes in snowpack are, therefore, expected to have direct, near-term influences on soil moisture profiles^{3,134} and ET fluxes. Anthropogenic warming enhances evaporative demand¹³⁵, which, subject to water availability, will increase ET. Yet, changing snowpacks can dramatically alter water availability, particularly during early spring and summer. Together, the increased evaporative demands along with changing water availability can make contemporary droughts similar in severity

to Medieval megadroughts^{136,137}. However, there are several mediating factors influencing and complicating snow change influences on ET, including: plant physiological changes^{138,139}, water-limitation caps to ET^{140,141}, vegetation access to water due to snowmelt timing shifts; and the degree to which subsurface storage can buffer changes in above ground snow. For example, assuming access to deep (>2 m) rooting zones and associated storage¹⁴², a warmer climate is anticipated to increase ET by 28% across a western Sierra Nevada watershed by the end of the century¹⁴³, with similar declines in streamflow. In contrast, other projections indicate reductions in ET by up to 40% for mid-latitude Sierra catchments, owing to earlier snowmelt and truncated water availability into the growing season¹⁴⁴. At longer, multi-year timescales, changes in the timing and magnitude of water availability for ET will both influence and respond to changes in the density and species composition of vegetation.

Groundwater. Groundwater recharge in mountainous systems, either diffuse (via direct percolation of precipitation into the subsurface) or focused (via surface water bodies), is expected to change, given alterations to snowpack¹⁴⁵. In general, snowmelt more effectively infiltrates into the subsurface when compared with rainfall¹⁴⁶. Thus, a phase change in future precipitation can result in less mountainous groundwater diffuse recharge. Furthermore, upward elevation shifts in the snowline will decrease the surface area over which snowmelt can occur⁷⁶, reducing snowmelt infiltration into lower permeability material and, thereby, decreasing overall recharge. If more snowpack becomes focused at higher elevations, there is potential for deeper snowpacks to accumulate, which melt later and faster¹²². However, low-permeability alpine snowmelt drives substantial groundwater interflow because ET demands in these regions are also low, enabling a substantial contribution of groundwater to subalpine vegetation — a potential climate-change-resilient mechanism for groundwater recharge¹⁴⁷. If changes in groundwater do occur, they can have cascading effects on subsurface weathering rates, chemical export¹⁴⁸ and discharge for groundwater gaining streams and rivers, with subsequent implications for groundwater-dependent ecosystems¹⁴⁹. There is, thus, a critical need to understand mountain groundwater in a future climate⁴⁰.

Runoff and streamflow. On average, snowmelt produces more runoff than rainfall^{115,150,151} and more attenuated peak flows than rainfall runoff¹⁵². Models suggest that more than half of runoff in the WUS originates from snow, despite snowfall representing only one-third of total precipitation. Snowmelt contributions to runoff in the mountains are even larger, up to 70%¹¹⁵, indicating that projected snowfall changes will influence streamflow.

Generally, a shift in precipitation phase from snow to rain leads to significant decreases in annual streamflow, reflecting increases in ET¹⁵⁰. However, as noted, the direct effects of a low-to-no snow future on ET in winter wet–summer dry systems can include both decreases and increases in ET. More water-limited areas might show decreases and higher elevation temperature-limited areas




Fig. 5 | Changes in mountain environments under persistent low-to-no snow conditions. a | Physical and biological processes as observed in mountain systems with seasonal snowpack. **b** | Shifts in the behaviour of mountain systems under future low-to-no snow. **c** | Seasonal trends of groundwater and streamflow throughout a single water year (beginning 1 October), where the blue line represents historical conditions and the red line represents hypothesized future conditions. Although predictions of many co-evolving hydrologic processes as a result of low-to-no snow conditions are difficult to infer, changes in both above and below ground future watershed behaviour is expected.

might show increases, and these changes will evolve with changes in vegetation distribution and type. Thus, while overall streamflow is likely to decrease, there could be locations where annual streamflow might increase with shifts from snow to rain. Yet, some projections reveal that mean future flows will remain stable, but that peak flows will increase and low flows will decrease¹⁵³. Changes to WUS hydrographs, however, should be contextualized relative to the low-to-no snow transition. For instance, in a warmer future where snowpack remains, it is expected that slower snowmelt rates will yield less streamflow¹⁵⁴. In a transitional future where both snow and rain exist, there is also increased potential for flood hazards due to rain-on-snow events^{77,155}. Rain-on-snow events effectively enhance runoff efficiency, where streamflow contributions from winter rainfall can surpass snowfall contributions¹⁵⁶.

At sub-annual timescales, seasonal low flow, fed predominately by late spring snowmelt and groundwater in gaining streams, is an important consideration, given its role in sustaining ecological habitats and water demands in Mediterranean climates, where annual peak water supply demands correspond to precipitation-free

months. Historical evaluation of snowpack and resulting streamflow relationships during low-discharge conditions reveals that, for every 10% decrease in peak SWE, annual minimum flows could decrease on the order of 9–22% and occur several days earlier¹⁵⁷. Shorter snowmelt seasons imply longer low-flow periods, which results in reduced low flows¹⁵⁸ (FIG. 5c). The translation of changes in the timing of snowmelt into low flows is strongly influenced by geological factors that influence watershed response times. Slower draining, deep groundwater-dominated watersheds such as the High Cascades in Oregon, for example, could show greater reduction in late-season flows relative to faster draining watersheds, such as steep granitic watersheds in the California Sierra Nevada or volcanic watersheds in the Western Oregon Cascades^{158,159}. As streamflow responses to changes in the magnitude and timing of snowmelt are strongly mediated by geologic factors that control recession characteristics, heterogeneity in geology, even within regions such as the California Sierra Nevada, will be an important consideration in hydrologic model applications to predict low-flow responses to changing snow¹⁵⁷.

Longer term and indirect hydrologic changes. At multi-year timescales, snowpack changes will impact the co-evolution of hydrologic and land-surface processes, such as the amount and type of vegetation present. For example, forest productivity in the California Sierra Nevada increases with snow accumulation, particularly for more water-limited mid-elevations¹⁶⁰, and earlier snowmelt reduces carbon uptake in the Colorado Rockies¹⁶¹, reflecting loss of water to support late summer productivity. Longer term species changes with warming are also expected¹⁶², along with increases in forest mortality with drought^{163,164} and disease^{165,166}. How changes in vegetation structure (including leaf area, rooting depth and biomass) and composition (water use access and water use efficiency) translate to changes in ET–streamflow partitioning remains poorly constrained in models, and is likely to vary with geoclimatic settings.

Changing vegetation structure can also feed back on the snowpack. For instance, denser forest canopies can increase snow interception losses through sublimation and increase snowmelt via greater longwave radiation. Denser canopies, however, can also slow snowmelt by reducing incoming shortwave radiation at the snow surface, and lessen sensible and latent heat fluxes into the sub-canopy snowpack¹⁶⁷. The relationship between forest cover and snow dynamics can also vary with local climate conditions and landscape morphology. Regionally, mean winter temperature is an indicator of where forest cover will act as an insulator to snow cover or enhance its melt; regions with December through February temperatures $>-1^{\circ}\text{C}$ reduce seasonal snow duration on the order of 1–2 weeks¹⁶⁸.

Changes in wildfire frequency, severity and timing are particularly catastrophic consequences of a low-to-no snow future. Indeed, alongside continued warming, a shift towards a no-snow future is anticipated to exacerbate wildfire activity, as observed^{169,170}. However, in the longer term, drier conditions can also slow post-fire vegetation regrowth, even reducing fire size and severity by reducing fuels. The hydrologic (and broader) impacts of fire are substantial, and include: shifts in snowpack accumulation, snowpack ablation and snowmelt timing¹⁷¹; increased probability of flash flooding and debris flows^{172,173}; enhanced overland flow; deleterious impacts on water quality^{174,175}; and increased sediment fluxes^{176,177}. Notably, even small increases in turbidity can directly impact water supply infrastructure^{178,179}. Vegetation recovery within the first few years following fire rapidly diminishes these effects, but some longer term effects do occur, as evidenced with stream chemistry¹⁸⁰ and above and below ground water partitioning both within and outside of burn scars¹⁸¹.

Integrated cascading responses. Projecting the propagation of changing snow dynamics onto ET, groundwater and surface water involves integrating cascading responses and their coupling with changing land-surface properties. Time horizons of these expected hydrologic and land-surface changes depend on the degree of low-to-no snow intermittency and other climate drivers, in turn, impacting the degree of system rebound. The potential consequences of different changes in annual to sub-seasonal

snowpack on major hydrologic stores and fluxes are substantial and difficult to infer, given simultaneous changes in temperature and precipitation (FIG. 6).

Regional-to-continental-scale estimates of the cascading effects of future low-snow scenarios on runoff are available for the WUS^{3,115}. However, the hydrologic models used for such analyses generally do not account for potential feedbacks related to changes in vegetation and land-surface characteristics, which are vital for longer range forecasts. Empirical analysis of hydrologic responses to wildfires¹⁸² show substantial regional variation within the WUS. Similarly, expected trajectories of changing wildfire regimes vary both within and between WUS regions¹⁸³. Coupled models must, therefore, resolve parameters and processes that drive not only significant spatial patterns in snow and hydrology but also spatial patterns of changing vegetation and its cascading impacts on water.


Hypothetical sequencing of low snow years can conceptually define transition times between conditions that are extreme, episodic or persistent (FIG. 6b), depending on their frequency and order. The occurrence of these transitions can subsequently impact the hydrologic dynamics. For example, groundwater response signals during high snow years can alleviate stresses of previous years during the extreme low-snow regimes, but not during episodic or persistent regimes (FIG. 6b). Hydrologic deviations from expected (or historic) responses will be dependent on the resilience of each component of the system, and the co-evolution and interaction of processes simultaneously. Thresholds in system response could then be reached more quickly, given disturbances (for example, fire or vegetation mortality induced by disease or insects).

Climate change adaptation strategies

In a future characterized by low-to-no snow, creative and flexible strategies to adapt to variable water supplies will be critical to reduce economic and ecosystem impacts, particularly in light of 71% of water experts believing that current water management strategies would be insufficient and the need for a better understanding of climate-change-induced uncertainties³³.

Vulnerabilities of the WUS water system. Over the last century, a complex engineered system has been developed in the WUS alongside the natural river network and snowpack storage to meet diverse water demands, including: rural, urban, residential, tribal, commercial, industrial, agricultural and hydroelectric power (FIG. 7). This system includes reservoirs for surface storage, canal and pump structures to move water long distances across basins and over mountains, groundwater pumps to augment surface supplies and hedge against dry years, run-of-river and reservoir-based hydropower to generate electricity, and water efficiency technologies and conservation practices to reduce water needs. In addition to ensuring reliable water supply, this system mitigates flood risk from high-water events and provides environmental flows for ecosystem benefits. Together, these systems have been designed to facilitate the movement of water from water-surplus to water-limited regions, to manage

a Single water year



b Multiple water years






Fig. 6 | Cascading above and below ground impacts of snowpack changes. a | Temporal dynamics of annual water partitioning for hypothetical variations (average snow year, low snow with earlier melt and low snow with rain on snow) in average and low snow years. With the exception of the rain-on-snow scenario, these illustrations assume no additional compensation from rain. Question marks denote scenarios where inferring potential changes are more uncertain. **b** | Sequencing effects of multi-year low-to-no snow conditions indicating transitions between extreme, episodic and persistent system behaviour. Red shading indicates low snow years and blue shading indicates high snow years. Inferences in system behaviour such as rebounds after single or sequential low snow years are difficult, given interactions across the atmosphere–bedrock continuum, warranting the use of physically based models. ET, evapotranspiration.

a Water sources, storage and conveyance



b Water demand





Fig. 7 | Supply and demand connections between the natural and managed WUS water systems. a | Water sources, storage and conveyance across the western United States (WUS). **b** | Water demands across the WUS. This combination of natural and engineered systems illustrates the interconnected nature of WUS water and the long travel distances of water moved away from mountains to support urban and agricultural water end users and to power run-of-river and reservoir-based hydropower generation. Note that groundwater aquifers, another major water source in the region, are not included in this figure for clarity of visual representation. See Supplementary Information for the data sources. SWE, snow water equivalent.

the natural spatiotemporal variability that is inherent in the WUS precipitation patterns and the locational and timing mismatch between water availability and human water demands across multiple sectors^{184,185}. The high value placed on water in these regions, particularly during times of scarcity, drives this movement of water across basins, despite the great financial and energy cost of doing so¹⁸⁶. A complicated set of legal commitments regulate much of the physical infrastructure, including long-term water compacts, interbasin transfers, water rights, water quality standards, and environmental water and species protection mandates. These physical and legal

structures are managed together in response to changing water demands across daily, yearly and decadal scales.

While WUS water management has adapted to inter-annual variability in water supply, fundamentally, the water infrastructure, legal framework of water rights¹⁸⁷ and management institutions were built with the expectation that supply variability would be contained within a stationary range of known possible conditions¹⁹. However, changes in the seasonality and variability of the mountainous hydrologic cycle has important direct and indirect implications for the management of WUS water resources that extend beyond just changes in

the historical mean and variability of annual runoff. For example, under a low-to-no snow future, it will be more challenging to capture earlier snowmelt in reservoirs required to maintain early season flood control capacity¹⁸⁸ or to gauge reservoir releases in order to prepare for secondary or tertiary peaks in inflow.

These changes will affect water availability for urban water suppliers, for example, in Southern California, a region heavily dependent on imported water from the snowpacks of the Sierra Nevada and Colorado River Basin¹⁸⁹. There will be increased pressure on agricultural water users in the WUS¹⁹⁰, who have come to rely on snowmelt stored in reservoirs to irrigate during the summer growing season^{6,191}. Rural community water systems, such as in the Central Valley of California, could become more reliant on already stressed and overdrafted groundwater sources, particularly those that already have water quality violations¹⁹². Climate change impacts, including declining snowpack, could create water supply availability and quality challenges in some tribal areas that exacerbate socio-economic vulnerabilities^{62,193,194}. In addition, a low-to-no snow future will affect the WUS energy sector; high-elevation hydropower generators that have small reservoirs designed around stationary snowpack storage will likely decrease their electricity production, particularly during the summer^{195–197}. Given that conveyance infrastructure connects disparate natural systems across hundreds of miles and between basins (FIG. 7), such impacts of low-to-no snow conditions in one region can have far-reaching impacts that disrupt the WUS water system across a diverse group of stakeholders with different economic values, water use priorities and legal mandates. These connections further compound existing pressures on water supplies from ageing infrastructure, population growth, increased demands from energy production and land-use change^{184,198}.

A path forward. A number of potential adaptation strategies to a low-to-no snow future can be pursued to address both water supply availability and water demands, with, for example, hard infrastructure (including dams, canals and new supplies) and soft management approaches (including better forecasting to inform operations, managed aquifer recharge (MAR) and conservation)¹⁹⁹. It is likely that a diversified portfolio of both supply-side and demand-side strategies will be needed to flexibly adapt to a low-to-no snow future, and overcome implementation barriers and physical limits of any one approach^{17,200}.

Physical surface storage and conveyance infrastructure (for example, dams, pipelines and aqueducts) serve to smooth the variability of water supplies in time and space. Proposed future adaptation strategies include building new surface storage capacity (for example, the Sites Reservoir in California)²⁰¹, retrofitting existing reservoirs to increase capacity (for example, raising Shasta Dam in California)^{202,203} and constructing large new conveyance projects (for example, Utah's proposed Lake Powell Pipeline on the Colorado River)²⁰⁴. However, dams and large conveyances are expensive, slow to develop relative to the speed of anthropogenic climate change and can have significant negative ecosystem

impacts, including obstructing and changing other physical aspects of fish habitats^{205,206}, trapping sediment²⁰⁷ and changing water quality. Planning of these new systems and associated water governance is also challenging because large infrastructure is designed, built and maintained on the multidecadal scale and includes trade-offs such as cost versus capacity, requiring decision frameworks and optimization paradigms that can account for multiple factors^{208,209}.

Besides new infrastructure, there is some flexibility to adapt existing surface storage infrastructure to new conditions through innovative reservoir operations. Forecast-informed reservoir operation (FIRO) is a relatively new concept in which weather and hydrologic forecasts are used to selectively retain or release water from reservoirs, as opposed to conventional release schedules based on rule curves, which are estimates of allowable maximum daily storage to reduce downstream flood risk^{84,210}. Advances in sub-seasonal to seasonal weather forecasting have made it possible for reservoir operators to more effectively balance water supply and flood control objectives (for example, avoid releasing water supply for flood preparedness if large precipitation events are unlikely in the near future). A modelling case study on Lake Mendocino showed the potential to increase water storage by 33%⁸⁴. Such strategies are invaluable as the need to capture spring snowmelt becomes less reliable in the water year. Enhanced monitoring networks can further support critical management decisions during these high-impact events²¹¹.

Groundwater reservoirs can also provide significant storage capacity to offset snow loss. For example, California groundwater basins are estimated to have between 850 million to 1.3 billion acre-feet (1,048 to 1,604 km³) of storage, compared with an annual average of 23.5 MAF or 29 km³ of surface water reservoir storage (with an upper bound of 50 MAF or 62 km³)^{5,212}. Therefore, MAR, whereby excess surface water is conveyed to flood-permeable landscapes to infiltrate into groundwater aquifers or stormwater is captured and stored in urban areas²¹³, could be another important climate adaptation strategy²¹⁴. Conveyance can transport water to locations suitable for groundwater recharge either at times when there is excess conveyance capacity or when coupled with FIRO to provide more storage space in reservoirs without 'losing' water from the managed system. Importantly, these potential new groundwater stores must be managed conjunctively, or in coordination with surface water, to allow critically overdrafted basins to recover and be maintained at more sustainable levels²¹⁵.

There are also potential approaches to increase total supply by bringing 'new' water into the system. These approaches can be natural-engineered hybrid approaches such as forest and vegetation management to reduce ET and increase snow storage, runoff and recharge²¹⁶. However, forest management for water could conflict with other desired forest management outcomes, and must consider a range of other objectives, including wildfire hazard, terrestrial carbon storage, habitat refugia, recreation impacts and economics of its scalability, again highlighting inherent trade-offs among

competing societal objectives^{217,218}. Cloud seeding is another hybrid system that has been pursued historically but carries with it controversy²¹⁹. Fully engineered approaches such as seawater and brackish desalination and recycled water for potable and non-potable reuse are also possibilities and face their own challenges in terms of cost, energy usage, social acceptance, implementation lags and capacity^{220–223}.

Water demand management can complement the aforementioned supply-side approaches. Agricultural water use can be reduced by improved irrigation efficiency, more precise water use versus yield estimates for crops, integrated crop water management, crop switching and land repurposing^{224–227}. Ideas for reducing water use in urban sectors include more water-efficient appliances, reduced outdoor irrigation, leak detection, rain-water capture and localized storage^{185,228}. Additionally, changes in agricultural and urban water markets and pricing could be powerful tools to manage demand and respond to scarcity²²⁹. All of these approaches could help to increase WUS water system resilience, but their implementation would be more optimal with an improved understanding of the spatiotemporal dynamics of a low-to-no snow future.

Decision-making under uncertainty and co-production.

Evaluating the risks and trade-offs among alternative climate adaptation strategies presents a scientific and policy challenge, given the highly variable, multiscale, interconnected and nonlinear nature of built infrastructure systems and the natural systems that they are embedded within. Upgrading water infrastructure so that it is prepared to respond to these types of changes and challenges will require enormous investment. For example, the American Society of Civil Engineers projects US\$1 trillion are needed to maintain and expand the nation's drinking water systems, which have received a grade of 'D' in their 2017 infrastructure ranking²³⁰. Determining which investments are worthwhile to pursue is a process that is informed by scientific and technical knowledge about hydroclimatic and infrastructure systems. For example, operational decision support models can be combined with economic²³¹ and bedrock through atmosphere modelling capabilities²³², to explore physically plausible future scenarios with consideration of mitigation strategies and water supply and demands. However, additional or improved scientific information is, by no means, the only barrier to adaptation^{233,234}, and related decision-making is fundamentally a social process that must resolve the trade-offs among multiple objectives arising from different communities, actors and institutions.

There is an extensive social science literature on climate adaptation that examines, for instance, institutional, cultural, behavioural and political aspects of decision-making^{235,236}, including which stakeholders have a voice in planning (often referred to as procedural justice)²³⁷ and who benefits from adaptations (often referred to as distributive justice)^{238,239}. There are several promising scientific developments that can serve to narrow uncertainties regarding a low-to-no snow future. These include real-time modelling and monitoring networks,

and the development of observationally informed coupled modelling strategies that represent key human and natural system processes in a self-consistent manner. While such advances in science can reduce uncertainty regarding future conditions, some degree of irreducible uncertainty will always remain²⁴⁰. Thus, at the same time that science evolves to increase predictive understanding of the mechanisms of hydroclimatic change, management practice must evolve to accommodate uncertainty regarding the changing patterns of current and future hydrologic variability. Developing a robust strategy and selecting investment options that balance competing societal objectives and multisectoral interactions (such as the interaction among water and energy¹⁸⁶ or water and carbon²⁰⁷ reduction goals) requires new approaches to integrate water resource planning. Frameworks and planning methods for decision-making under deep uncertainty that acknowledge and accommodate imperfect knowledge regarding the probabilistic range of possible future conditions such as decision scaling²⁴¹, robust decision-making, dynamic adaptation pathways²⁴² and scenario planning can identify scientifically informed adaptive strategies that leverage best available science without overstating its confidence²⁴³.

For instance, the United States Bureau of Reclamation and water management agencies within the Colorado River Basin engaged in a robust decision-making study that identified a range of potential future climate conditions under which water delivery obligations would be vulnerable. Portfolios of adaptation strategies aimed at demand reduction (including agricultural, municipal and industrial conservation) and supply augmentation (including reuse, desalination and water import) were evaluated for their ability to alleviate these vulnerabilities and for their trade-offs in cost, yield, technical feasibility, legal risk and other criteria. The portfolios generally increase system robustness but have a wide range of implementation costs, especially under the declining supply conditions, and vary between the Upper Basin and the Lower Basin²⁴⁴.

Making science usable for decision-making requires strong trust between the parties²⁴⁵. This trust often develops over deliberate, long-term collaboration²⁴⁶, with mutual understanding of the science, models and tools being discussed and demonstration of the credibility, saliency and legitimacy of the new approach(es)²⁴⁷. Institutional, technical and financial capacity to implement these approaches must also be overcome²³³. Scientists must also recognize that practitioners are often directly responsible, sometimes even personally liable, for the outcomes of decisions made, which makes them hesitant in the application of new climate science²³⁶, especially if perceived as not fitting with existing knowledge or policy goals^{233,248}.

A path forward can be made by including Earth scientists, infrastructure experts, decision scientists, water management practitioners and community stakeholders, in a collaborative, iterative process of scientific knowledge creation through a co-production framework^{41,42,249,250}. This process helps to ensure that new science is suited to challenges at hand and can provide meaningful input into decision-making processes,

enabling more efficient, flexible and robust solutions. For example, an interdisciplinary, social and hydroclimatic co-production project between the Eastern Shoshone and Northern Arapaho tribes of the Wind River Indian Reservation in Wyoming, university partners and more than a dozen government agencies helped to develop early-warning tools for drought and to support climate adaptation planning at scales relevant to local water managers, in response to climate information needs identified by the tribes⁶². In such collaborations, boundary spanners or information brokers, organizations and/or individuals who can facilitate the co-production process play critical convening and translating roles in increasing the usability of new climate science for adaptation planning^{234,248,251,252}.

Summary and future perspectives

The WUS water sector built its infrastructure and developed its management practices under the assumption of an abundant winter snowpack that reliably melts prior to peak water demands in summer months. Observations highlight multiple lines of evidence indicating a historically declining snowpack that peaks and melts earlier. Models further project a continuation of this decline, with equivalent snowpack losses on the order of $\sim 35 \pm 10\%$ by mid-century and $\sim 50 \pm 10\%$ before the end of the twenty-first century, should emissions continue unabated. Exact timelines of snowpack declines are uncertain, shaped by projection methods, emission scenario and metrics used. To standardize community efforts to understand the time to emergence of snow loss, a new low-to-no snow definition is provided based on historical peak SWE percentiles, ≤ 30 th percentile (low snow) and ≤ 10 th percentile (effectively no snow). This low-to-no snow definition helps characterize when snow loss becomes particularly detrimental, which is likely regionally specific and will be a function of the sequencing of low snow years and the response of the hydrologic system to those low snow years. The implications of these snowpack changes include perturbations to groundwater recharge, streamflow dynamics and ET fluxes, and are complicated by simultaneous, nonlinear processes from the bedrock through the canopy, as well as disturbance such as wildfire. Owing to the expansive and interconnected nature of WUS water storage and conveyance, particularly to differential snow loss in various mountain regions, several vulnerabilities to the existing WUS water system will require innovative adaptation approaches that can be supported by new modes of usable, decision-relevant knowledge production.

Of the many uncertainties imposed by climate change on the hydrologic cycle, the high confidence in snowpack decline enables proactive planning. From a water storage perspective alone, a shift in precipitation from snow to rain will result in a failure to meet water demands, given limitations in existing reservoir storage capacity. Strategies could include additional surface and subsurface storage, diversifying supply options and conservation. A re-evaluation of the operation and design of WUS water systems is timely, given current plans for large-scale federal investments in infrastructure

and the American Society of Civil Engineers' recent declaration of 'chronically underinvested' US drinking water systems²³⁰. This re-evaluation also presents an opportunity to proactively plan for snow-related water failures. Snow loss is often less of a priority relative to other investments made retroactively to a changing climate, for example, Texas' winterization from the national electric grid or the US coastal flood defence from storm surges and sea level rise²⁵³. For water management, the urgency of implementation for the planning, permitting, construction and/or retrofitting of storage and conveyance systems is immediate, but the realization of benefits is on the order of several decades¹⁸⁴. Thus, decisions and investments made today will extend multiple generations, operate for half-centuries or more, and need to function within rapidly changing hydroclimatic conditions. To avoid pitfalls of assuming stationarity again, investments should not solely focus on hard infrastructure but also the potential for soft management approaches and continuous optimization of the hard infrastructure investments, such as MAR and FIRO, as environmental conditions change.

In combination with hard infrastructure and soft management investments, additional investments in maintaining and expanding observational networks and supporting the implementation of frameworks to develop co-produced, use-inspired research will be instrumental as snow disappears. Gaps between operational and research models need to be addressed. Future ESM development aimed at enhancing the fidelity of integrated mountainous hydrologic cycle processes should also be considered a scientific grand challenge. The overall goal being that future models that can reliably translate projections of planetary-to-regional-scale changes in the Earth's climate system to the scales relevant to the mountainous hydrologic cycle is, in turn, better suited to isolate existing natural and managed system vulnerabilities and assess the efficacy and scalability of climate adaptation strategies.

Finally, a paradigm shift in the science community is also needed, where those studying climate change will need to not only look for a single mode of hypothesis-driven research but also towards a problem-oriented framework²⁵⁴. Likewise, water agencies will need to be amenable to new technologies, approaches and develop non-traditional relationships to inform new protocols for operations. This more holistic approach that accepts nonstationarity while considering both the natural physical system and the managed one can help to ensure that systems are less vulnerable to longer term hydroclimatic change and short-term hydrometeorological extremes. A concerted, community-wide effort is needed to support the funding mechanisms, legal and research institutions, incentive structures, co-production teams and/or boundary organizations to represent a diverse set of stakeholder needs. The overall goal of these investments and interactions would be WUS-wide resilience to ongoing snow loss and the imminent onset of a persistent low-to-no snow future if anthropogenic climate change continues unabated.

Published online 26 October 2021

1. Huss, M. et al. Toward mountains without permanent snow and ice. *Earths Future* **5**, 418–435 (2017).
2. Immerzeel, W. W. et al. Importance and vulnerability of the world's water towers. *Nature* **577**, 364–369 (2020).
3. Gergel, D. R., Nijssen, B., Abatzoglou, J. T., Lettenmaier, D. P. & Stumbaugh, M. R. Effects of climate change on snowpack and fire potential in the western USA. *Clim. Change* **141**, 287–299 (2017).
4. Sturm, M., Goldstein, M. A. & Parr, C. Water and life from snow: a trillion dollar science question. *Water Resour. Res.* **53**, 3534–3544 (2017).
5. Dettinger, M. D. & Anderson, M. L. Storage in California's reservoirs and snowpack in this time of drought. *San Franc. Estuary Watershed Sci.* **13**, 1 (2015).
6. Qin, Y. et al. Agricultural risks from changing snowmelt. *Nat. Clim. Chang.* **10**, 459–465 (2020).
7. Rhoades, A. M., Ullrich, P. A. & Zarzycki, C. M. Projecting 21st century snowpack trends in western USA mountains using variable-resolution CESM. *Clim. Dyn.* **50**, 261–288 (2018).
8. Pierce, D. W. et al. Probabilistic estimates of future changes in California temperature and precipitation using statistical and dynamical downscaling. *Clim. Dyn.* **40**, 839–856 (2013).
9. Marshall, A. M., Abatzoglou, J. T., Link, T. E. & Tennant, C. J. Projected changes in interannual variability of peak snowpack amount and timing in the Western United States. *Geophys. Res. Lett.* **46**, 8882–8892 (2019).
10. Barnett, T. P., Adam, J. C. & Lettenmaier, D. P. Potential impacts of a warming climate on water availability in snow-dominated regions. *Nature* **438**, 303–309 (2005).
11. Allan, R. P. et al. Advances in understanding large-scale responses of the water cycle to climate change. *Ann. NY Acad. Sci.* **1472**, 49–75 (2020).
12. Huning, L. S. & AghaKouchak, A. Global snow drought hot spots and characteristics. *Proc. Natl Acad. Sci. USA* **117**, 19755–19759 (2020).
13. Polade, S. D., Gershunov, A., Cayan, D. R., Dettinger, M. D. & Pierce, D. W. Precipitation in a warming world: assessing projected hydro-climate changes in California and other Mediterranean climate regions. *Sci. Rep.* **7**, 10783 (2017).
14. Milly, P. C. D. & Dunne, K. A. Colorado River flow dwindles as warming-driven loss of reflective snow energizes evaporation. *Science* **367**, 1252–1255 (2020).
15. Mote, P. W., Li, S., Lettenmaier, D. P., Xiao, M. & Engel, R. Dramatic declines in snowpack in the western US. *npj Clim. Atmos. Sci.* **1**, 2 (2018).
16. Mountain Research Initiative EDW Working Group. Elevation-dependent warming in mountain regions of the world. *Nat. Clim. Chang.* **5**, 424–430 (2015).
17. US Global Change Research Program. Climate science special report: fourth national climate assessment, Vol. I (USGCRP, 2017).
18. Rhoades, A. M., Jones, A. D. & Ullrich, P. A. Assessing mountains as natural reservoirs with a multimetric framework. *Earths Future* **6**, 1221–1241 (2018).
19. Milly, P. C. D. et al. Stationarity is dead: whither water management? *Earth* **4**, 20 (2008).
20. Vano, J. A. Implications of losing snowpack. *Nat. Clim. Chang.* **10**, 388–390 (2020).
21. Ullrich, P. A. et al. California's drought of the future: a midcentury recreation of the exceptional conditions of 2012–2017. *Earths Future* **6**, 1568–1587 (2018).
22. Livneh, B. & Badger, A. M. Drought less predictable under declining future snowpack. *Nat. Clim. Chang.* **10**, 452–458 (2020).
23. US Geophysics Research Forum & Geophysics Study Committee. *Scientific Basis of Water-resource Management* (National Academy, 1982).
24. Hatchett, B. J., Boyle, D. P., Putnam, A. E. & Bassett, S. D. Placing the 2012–2015 California-Nevada drought into a paleoclimatic context: Insights from Walker Lake, California-Nevada, USA. *Geophys. Res. Lett.* **42**, 8632–8640 (2015).
25. Hatchett, B. J. et al. in *From Saline to Freshwater: The Diversity of Western Lakes in Space and Time*, Geological Society of America Special Paper Vol. 536 (eds Starratt, S. W. & Rosen, M. R.) 67–79 (Geological Society of America, 2019).
26. Woodhouse, C. A., Kunkel, K. E., Easterling, D. R. & Cook, E. R. The twentieth-century pluvial in the western United States. *Geophys. Res. Lett.* **32**, L07701 (2005).
27. Cook, B. I., Seager, R. & Miller, R. L. On the causes and dynamics of the early twentieth-century North American pluvial. *J. Clim.* **24**, 5043–5060 (2011).
28. Kreuzt, K. J. et al. Bipolar changes in atmospheric circulation during the Little Ice Age. *Science* **277**, 1294–1296 (1997).
29. Hatchett, B. J. & McEvoy, D. J. Exploring the origins of snow drought in the northern Sierra Nevada, California. *Earth Interact.* **22**, 1–13 (2018).
30. Noble, P. J. et al. Holocene paleoclimate history of Fallen Leaf Lake, CA., from geochemistry and sedimentology of well-dated sediment cores. *Quat. Sci. Rev.* **131**, 193–210 (2016).
31. Shuman, B. N. & Serravezza, M. Patterns of hydroclimatic change in the Rocky Mountains and surrounding regions since the last glacial maximum. *Quat. Sci. Rev.* **173**, 58–77 (2017).
32. Bacon, S. N., Lancaster, N., Stine, S., Rhodes, E. J. & McCarley Holder, G. A. A continuous 4000-year lake-level record of Owens Lake, south-central Sierra Nevada, California, USA. *Quat. Res.* **90**, 276–302 (2018).
33. Sterle, K. & Singletary, L. Adapting to variable water supply in the Truckee-Carson river system, western USA. *Water* **9**, 768 (2017).
34. Hossain, F. et al. What do experienced water managers think of water resources of our nation and its management infrastructure? *PLoS ONE* **10**, e0142073 (2015).
35. Lynn, E. et al. Technical note: Precipitation-phase partitioning at landscape scales to regional scales. *Hydrol. Earth Syst. Sci.* **24**, 5317–5328 (2020).
36. Nolin, A. W. & Daly, C. Mapping "at risk" snow in the Pacific Northwest. *J. Hydrometeorol.* **7**, 1164–1171 (2006).
37. Mott, R., Vionnet, V. & Grünwald, T. The seasonal snow cover dynamics: review on wind-driven coupling processes. *Front. Earth Sci.* **6**, 197 (2018).
38. Shafer, S. L., Bartlein, P. J. & Whitlock, C. in *Global Change and Mountain Regions: An Overview of Current Knowledge* (eds Huber, U. M., Bugmann, H. K. M. & Reasoner, M. A.) 21–30 (Springer, 2005).
39. Maina, F. Z., Siirila-Woodburn, E. R., Newcomer, M., Xu, Z. & Steefel, C. Determining the impact of a severe dry to wet transition on watershed hydrodynamics in California, USA with an integrated hydrologic model. *J. Hydrol.* **580**, 124358 (2020).
40. Meixner, T. et al. Implications of projected climate change for groundwater recharge in the western United States. *J. Hydrol.* **534**, 124–138 (2016).
41. Jagannathan, K. et al. Great expectations? Reconciling the aspiration, outcome, and possibility of co-production. *Curr. Opin. Environ. Sustain.* **42**, 22–29 (2020).
42. Jagannathan, K., Jones, A. D. & Ray, I. The making of a metric: co-producing decision-relevant climate science. *Bull. Am. Meteorol. Soc.* **102**, E1579–E1590 (2021).
43. Neelin, J. D., Langenbrunner, B., Meyerson, J. E., Hall, A. & Berg, N. California winter precipitation change under global warming in the Coupled Model Intercomparison Project phase 5 ensemble. *J. Clim.* **26**, 6238–6256 (2013).
44. Dong, L., Leung, L. R., Lu, J. & Song, F. Double-ITCZ as an emergent constraint for future precipitation over Mediterranean climate regions in the North Hemisphere. *Geophys. Res. Lett.* **48**, e2020GL091569 (2021).
45. Persad, G. G., Swain, D. L., Kouba, C. & Ortiz-Partida, J. P. Inter-model agreement on projected shifts in California hydroclimate characteristics critical to water management. *Clim. Change* **162**, 1493–1513 (2020).
46. Helms, D., Phillips, S. E. & Reich, P. F. *The History of Snow Survey and Water Supply Forecasting: Interviews with U.S. Department of Agriculture Pioneers* (US Department of Agriculture, 2008).
47. Huning, L. S. & AghaKouchak, A. Approaching 80 years of snow water equivalent information by merging different data streams. *Sci. Data* **7**, 333 (2020).
48. Evan, A. T. A new method to characterize changes in the seasonal cycle of snowpack. *J. Appl. Meteorol. Climatol.* **58**, 131–143 (2019).
49. Margulis, S. A., Liu, Y. & Baldo, E. A joint landsat- and MODIS-based reanalysis approach for midlatitude montane seasonal snow characterization. *Front. Earth Sci.* **7**, 272 (2019).
50. Kapnick, S. & Hall, A. Causes of recent changes in western North American snowpack. *Clim. Dyn.* **38**, 1885–1899 (2012).
51. Kapnick, S. & Hall, A. Observed climate–snowpack relationships in California and their implications for the future. *J. Clim.* **23**, 3446–3456 (2010).
52. Margulis, S. A. et al. Characterizing the extreme 2015 snowpack deficit in the Sierra Nevada (USA) and the implications for drought recovery. *Geophys. Res. Lett.* **43**, 6341–6349 (2016).
53. Margulis, S. A., Cortés, G., Giroto, M. & Durand, M. A Landsat-era Sierra Nevada snow reanalysis (1985–2015). *J. Hydrometeorol.* **17**, 1203–1221 (2016).
54. Hatchett, B. J. Seasonal and ephemeral snowpacks of the conterminous United States. *Hydrology* **8**, 32 (2021).
55. Bohr, G. S. & Aguado, E. Use of April 1 SWE measurements as estimates of peak seasonal snowpack and total cold-season precipitation. *Water Resour. Res.* **37**, 51–60 (2001).
56. Hatchett, B. J., Rhoades, A. M. & McEvoy, D. J. Monitoring the daily evolution and extent of snow drought. Preprint at *Nat. Hazards Earth Syst. Sci.* <https://doi.org/10.5194/nhess-2021-193> (2021).
57. Harpold, A., Dettinger, M. & Rajagopal, S. Defining snow drought and why it matters. *Eos* <https://doi.org/10.1029/2017eo068775> (2017).
58. Dierauer, J. R., Allen, D. M. & Whitfield, P. H. Snow drought risk and susceptibility in the western United States and southwestern Canada. *Water Resour. Res.* **55**, 3076–3091 (2019).
59. Rasmussen, R. et al. How well are we measuring snow: the NOAA/FAA/NCAR winter precipitation test bed. *Bull. Am. Meteorol. Soc.* **93**, 811–829 (2012).
60. Kinar, N. J. & Pomeroy, J. W. Measurement of the physical properties of the snowpack. *Rev. Geophys.* **53**, 481–544 (2015).
61. Lundquist, J., Hughes, M., Gutmann, E. & Kapnick, S. Our skill in modeling mountain rain and snow is bypassing the skill of our observational networks. *Bull. Am. Meteorol. Soc.* **100**, 2473–2490 (2019).
62. McNeely, S. M. et al. Anatomy of an interrupted irrigation season: micro-drought at the Wind River Indian Reservation. *Clim. Risk Manag.* **19**, 61–82 (2018).
63. Huning, L. S. & AghaKouchak, A. Mountain snowpack response to different levels of warming. *Proc. Natl Acad. Sci. USA* **115**, 10932–10937 (2018).
64. Wrzesien, M. L., Paveisky, T. M., Durand, M. T., Dozier, J. & Lundquist, J. D. Characterizing biases in mountain snow accumulation from global data sets. *Water Resour. Res.* **55**, 9873–9891 (2019).
65. Painter, T. H. et al. The Airborne Snow Observatory: fusion of scanning lidar, imaging spectrometer, and physically-based modeling for mapping snow water equivalent and snow albedo. *Remote Sens. Environ.* **184**, 139–152 (2016).
66. Guan, B. et al. Snow water equivalent in the Sierra Nevada: blending snow sensor observations with snowmelt model simulations. *Water Resour. Res.* **49**, 5029–5046 (2013).
67. Rittger, K., Bair, E. H., Kahl, A. & Dozier, J. Spatial estimates of snow water equivalent from reconstruction. *Adv. Water Resour.* **94**, 345–363 (2016).
68. Crumley, R. L. et al. Assimilation of citizen science data in snowpack modeling using a new snow dataset: Community Snow Observations. *Hydrol. Earth Syst. Sci.* **25**, 4651–4680 (2021).
69. Musselman, K. N., Addor, N., Vano, J. A. & Molotch, N. P. Winter melt trends portend widespread declines in snow water resources. *Nat. Clim. Chang.* **1**, 418–424 (2021).
70. Harpold, A. et al. Changes in snowpack accumulation and ablation in the intermountain west. *Water Resour. Res.* **48**, W11501 (2012).
71. Mote, P. W., Hamlet, A. F. & Clark, M. P. Declining mountain snowpack in western North America. *Bull. Am. Meteorol. Soc.* **86**, 39–50 (2005).
72. Michelle Hu, J. & Nolin, A. W. Widespread warming trends in storm temperatures and snowpack fate across the Western United States. *Environ. Res. Lett.* **15**, 034059 (2020).
73. Gonzales, K. R., Swain, D. L., Nardi, K. M., Barnes, E. A. & Diffenbaugh, N. S. Recent warming of landfalling atmospheric rivers along the west coast of the United States. *J. Geophys. Res.* **24**, 6810–6826 (2019).
74. Payne, A. E. et al. Responses and impacts of atmospheric rivers to climate change. *Nat. Rev. Earth Environ.* **1**, 143–157 (2020).
75. Oyler, J. W., Dobrowski, S. Z., Ballantyne, A. P., Klene, A. E. & Running, S. W. Artificial amplification of warming trends across the mountains of the western United States. *Geophys. Res. Lett.* **42**, 153–161 (2015).
76. Wayand, N. E., Lundquist, J. D. & Clark, M. P. Modeling the influence of hypsometry, vegetation,

- and storm energy on snowmelt contributions to basins during rain-on-snow floods. *Water Resour. Res.* **51**, 8551–8569 (2015).
77. McCabe, G. J., Clark, M. P. & Hay, L. E. Rain-on-snow events in the western United States. *Bull. Am. Meteorol. Soc.* **88**, 319–328 (2007).
 78. Skiles, S. M., Flanner, M., Cook, J. M., Dumont, M. & Painter, T. H. Radiative forcing by light-absorbing particles in snow. *Nat. Clim. Chang.* **8**, 964–971 (2018).
 79. Gleason, K. E., McConnell, J. R., Arienzo, M. M., Chellman, N. & Calvin, W. M. Four-fold increase in solar forcing on snow in western US burned forests since 1999. *Nat. Commun.* **10**, 2026 (2019).
 80. AghaKouchak, A. et al. How do natural hazards cascade to cause disasters? *Nature* **561**, 458–460 (2018).
 81. Parks, S. A. & Abatzoglou, J. T. Warmer and drier fire seasons contribute to increases in area burned at high severity in western US forests from 1985 to 2017. *Geophys. Res. Lett.* **47**, e2020GL089858 (2020).
 82. Neff, J. C. et al. Increasing eolian dust deposition in the western United States linked to human activity. *Nat. Geosci.* **1**, 189–195 (2008).
 83. Skiles, S. M. et al. Implications of a shrinking Great Salt Lake for dust on snow deposition in the Wasatch Mountains, UT, as informed by a source to sink case study from the 13–14 April 2017 dust event. *Environ. Res. Lett.* **13**, 124031 (2018).
 84. Delaney, C. J. et al. Forecast informed reservoir operations using ensemble streamflow predictions for a multipurpose reservoir in northern California. *Water Resour. Res.* **56**, e2019WR026604 (2020).
 85. Roberts, M., Milman, A. & Blomquist, W. in *Water Resilience: Management and Governance in Times of Change* (eds Baird, J. & Plummer, R.) 41–63 (Springer, 2021).
 86. Maina, F. Z., Siirila-Woodburn, E. R. & Vahmani, P. Sensitivity of meteorological-forcing resolution on hydrologic variables. *Hydrol. Earth Syst. Sci.* **24**, 3451–3474 (2020).
 87. Randall, D. A. et al. 100 years of earth system model development. *Meteorol. Monogr.* **59**, 12.1–12.66 (2018).
 88. Leutwyler, D., Fuhrer, O., Lapillonne, X., Lüthi, D. & Schär, C. Towards European-scale convection-resolving climate simulations with GPUs: a study with COSMO 4.19. *Geosci. Model Dev.* **9**, 3393–3412 (2016).
 89. Schär, C. et al. Kilometer-scale climate models: prospects and challenges. *Bull. Am. Meteorol. Soc.* **101**, E567–E587 (2020).
 90. Satoh, M. et al. Global cloud-resolving models. *Curr. Clim. Change Rep.* **5**, 172–184 (2019).
 91. Martin Ralph, F. et al. West Coast forecast challenges and development of atmospheric river reconnaissance. *Bull. Am. Meteorol. Soc.* **101**, E1357–E1377 (2020).
 92. Ralph, F. M. et al. A vision for future observations for western U.S. extreme precipitation and flooding. *J. Contemp. Water Res. Educ.* **153**, 16–32 (2014).
 93. Hatchett, B. J. et al. Observations of an extreme atmospheric river storm with a diverse sensor network. *Earth Space Sci.* **7**, e2020EA001129 (2020).
 94. White, A. B. et al. A twenty-first-century California observing network for monitoring extreme weather events. *J. Atmos. Ocean. Technol.* **30**, 1585–1603 (2013).
 95. Shen, C. A transdisciplinary review of deep learning research and its relevance for water resources scientists. *Water Resour. Res.* **54**, 8558–8593 (2018).
 96. Schneider, T., Lan, S., Stuart, A. & Teixeira, J. Earth system modeling 2.0: a blueprint for models that learn from observations and targeted high-resolution simulations. *Geophys. Res. Lett.* **44**, 12,396–12,417 (2017).
 97. Yang, Q. et al. Exploring the effects of a nonhydrostatic dynamical core in high-resolution aquaplanet simulations. *J. Geophys. Res.* **122**, 3245–3265 (2017).
 98. Arakawa, A. & Jung, J.-H. Multiscale modeling of the moist-convective atmosphere — A review. *Atmos. Res.* **102**, 263–285 (2011).
 99. Gross, M. et al. Physics–dynamics coupling in weather, climate, and earth system models: challenges and recent progress. *Mon. Weather Rev.* **146**, 3505–3544 (2018).
 100. Harris, L. M., Lin, S.-J. & Tu, C. High-resolution climate simulations using GFDL HiRAM with a stretched global grid. *J. Clim.* **29**, 4293–4314 (2016).
 101. Zarzycki, C. M., Jablonowski, C. & Taylor, M. A. Using variable-resolution meshes to model tropical cyclones in the community atmosphere model. *Mon. Weather Rev.* **142**, 1221–1239 (2014).
 102. Rauscher, S. A. & Ringler, T. D. Impact of variable-resolution meshes on midlatitude baroclinic eddies using CAM-MPAS-A. *Mon. Weather Rev.* **142**, 4256–4268 (2014).
 103. Rhoades, A. M., Huang, X., Ullrich, P. A. & Zarzycki, C. M. Characterizing Sierra Nevada snowpack using variable-resolution CESM. *J. Appl. Meteorol. Climatol.* **55**, 173–196 (2016).
 104. Ghan, S. J. & Shippert, T. Physically based global downscaling: climate change projections for a full century. *J. Clim.* **19**, 1589–1604 (2006).
 105. Qian, Y., Ghan, S. J. & Leung, L. R. Downscaling hydroclimatic changes over the Western US based on CAM subgrid scheme and WRF regional climate simulations. *Int. J. Climatol.* **30**, 675–693 (2009).
 106. Clark, M. P. et al. A unified approach for process-based hydrologic modeling: 2. Model implementation and case studies. *Water Resour. Res.* **51**, 2515–2542 (2015).
 107. Fisher, R. A. & Koven, C. D. Perspectives on the future of land surface models and the challenges of representing complex terrestrial systems. *J. Adv. Model. Earth Syst.* **12**, e2018MS001453 (2020).
 108. Ning, L., Zhan, C., Luo, Y., Wang, Y. & Liu, L. A review of fully coupled atmosphere-hydrology simulations. *J. Geogr. Sci.* **29**, 465–479 (2019).
 109. Smyth, E. J., Raleigh, M. S. & Small, E. E. Improving SWE estimation with data assimilation: the influence of snow depth observation timing and uncertainty. *Water Resour. Res.* **56**, e2019WR026853 (2020).
 110. Margulis, S. A., Fang, Y., Li, D., Lettenmaier, D. P. & Andreadis, K. The utility of infrequent snow depth images for deriving continuous space-time estimates of seasonal snow water equivalent. *Geophys. Res. Lett.* **46**, 5331–5340 (2019).
 111. Krinner, G. et al. ESM-SnowMIP: assessing snow models and quantifying snow-related climate feedbacks. *Geosci. Model Dev.* **11**, 5027–5049 (2018).
 112. Rhoades, A. M., Jones, A. D. & Ullrich, P. A. The changing character of the California Sierra Nevada as a natural reservoir. *Geophys. Res. Lett.* **45**, 13,008–13,019 (2018).
 113. Pierce, D. W. & Cayan, D. R. The uneven response of different snow measures to human-induced climate warming. *J. Clim.* **26**, 4148–4167 (2013).
 114. Fyfe, J. C. et al. Large near-term projected snowpack loss over the western United States. *Nat. Commun.* **8**, 14996 (2017).
 115. Li, D., Wrzesien, M. L., Durand, M., Adam, J. & Lettenmaier, D. P. How much runoff originates as snow in the western United States, and how will that change in the future? *Geophys. Res. Lett.* **44**, 6163–6172 (2017).
 116. López-Moreno, J. I. et al. Different sensitivities of snowpacks to warming in Mediterranean climate mountain areas. *Environ. Res. Lett.* **12**, 074006 (2017).
 117. Rasmussen, R. et al. Climate change impacts on the water balance of the Colorado headwaters: high-resolution regional climate model simulations. *J. Hydrometeorol.* **15**, 1091–1116 (2014).
 118. Alder, J. R. & Hostetler, S. W. The dependence of hydroclimate projections in snow-dominated regions of the western United States on the choice of statistically downscaled climate data. *Water Resour. Res.* **55**, 2279–2300 (2019).
 119. Leung, L. R. et al. Mid-century ensemble regional climate change scenarios for the western United States. *Clim. Change* **62**, 75–113 (2004).
 120. Wrzesien, M. L. & Pavelsky, T. M. Projected changes to extreme runoff and precipitation events from a downscaled simulation over the western United States. *Front. Earth Sci.* **7**, 355 (2020).
 121. Mahoney, K. et al. Cool season precipitation projections for California and the Western United States in NA-CORDEX models. *Clim. Dyn.* **56**, 3081–3102 (2021).
 122. Musselman, K. N., Clark, M. P., Liu, C., Ikeda, K. & Rasmussen, R. Slower snowmelt in a warmer world. *Nat. Clim. Chang.* **7**, 214–219 (2017).
 123. McCrary, R. & Mearns, L. O. Quantifying and diagnosing sources of uncertainty in midcentury changes in North American snowpack from NARCCAP. *J. Hydrometeorol.* **20**, 2229–2252 (2019).
 124. Berg, N. & Hall, A. Anthropogenic warming impacts on California snowpack during drought. *Geophys. Res. Lett.* **44**, 2511–2518 (2017).
 125. Gutowski, W. J. et al. The ongoing need for high-resolution regional climate models: process understanding and stakeholder information. *Bull. Am. Meteorol. Soc.* **101**, E664–E683 (2020).
 126. Morss, R. E., Wilhelmi, O. V., Downton, M. W. & Grunfest, E. Flood risk, uncertainty, and scientific information for decision making: lessons from an interdisciplinary project. *Bull. Am. Meteorol. Soc.* **86**, 1593–1602 (2005).
 127. Svoboda, M. et al. The drought monitor. *Bull. Am. Meteorol. Soc.* **83**, 1181–1190 (2002).
 128. Lehner, F. et al. Partitioning climate projection uncertainty with multiple large ensembles and CMIP5/6. *Earth Syst. Dyn.* **11**, 491–508 (2020).
 129. Deser, C. et al. Insights from Earth system model initial-condition large ensembles and future prospects. *Nat. Clim. Chang.* **10**, 277–286 (2020).
 130. Hausfather, Z. & Peters, G. P. Emissions—the ‘business as usual’ story is misleading. *Nature* **577**, 618–620 (2020).
 131. Mizuta, R. et al. Climate simulations using MRI-AGCM3.2 with 20-km grid. *J. Meteorol. Soc. Japan* **90A**, 233–258 (2012).
 132. Haarsma, R. J. et al. High resolution model intercomparison project (HighResMIP v1.0) for CMIP6. *Geosci. Model Dev.* **9**, 4185–4208 (2016).
 133. Siler, N., Proistosescu, C. & Po-Chedley, S. Natural variability has slowed the decline in western U.S. snowpack since the 1980s. *Geophys. Res. Lett.* **46**, 346–355 (2019).
 134. Hamlet, A. F., Mote, P. W., Clark, M. P. & Lettenmaier, D. P. Twentieth-century trends in runoff, evapotranspiration, and soil moisture in the western United States. *J. Clim.* **20**, 1468–1486 (2007).
 135. Diffenbaugh, N. S., Swain, D. L. & Touma, D. Anthropogenic warming has increased drought risk in California. *Proc. Natl Acad. Sci. USA* **112**, 3931–3936 (2015).
 136. Woodhouse, C. A., Meko, D. M., MacDonald, G. M., Stahle, D. W. & Cook, E. R. A 1,200-year perspective of 21st century drought in southwestern North America. *Proc. Natl Acad. Sci. USA* **107**, 21283–21288 (2010).
 137. Stine, S. Extreme and persistent drought in California and Patagonia during mediaeval time. *Nature* **369**, 546–549 (1994).
 138. Katul, G. G., Oren, R., Manzoni, S., Higgins, C. & Parlange, M. B. Evapotranspiration: a process driving mass transport and energy exchange in the soil-plant-atmosphere-climate system. *Rev. Geophys.* **50**, RG3002 (2012).
 139. Roderick, M. L., Greve, P. & Farquhar, G. D. On the assessment of aridity with changes in atmospheric CO₂. *Water Resour. Res.* **51**, 5450–5463 (2015).
 140. Condon, L. E., Atchley, A. L. & Maxwell, R. M. Evapotranspiration depletes groundwater under warming over the contiguous United States. *Nat. Commun.* **11**, 873 (2020).
 141. Jung, M. et al. Recent decline in the global land evapotranspiration trend due to limited moisture supply. *Nature* **467**, 951–954 (2010).
 142. Klos, P. Z. et al. Subsurface plant-accessible water in mountain ecosystems with a Mediterranean climate. *Wiley Interdiscip. Rev. Water* **5**, e1277 (2018).
 143. Goulden, M. L. & Bales, R. C. Vulnerability of montane runoff to increased evapotranspiration with upslope vegetation distribution. *Proc. Natl Acad. Sci. USA* **111**, 14071–14075 (2014).
 144. Tague, C. & Peng, H. The sensitivity of forest water use to the timing of precipitation and snowmelt recharge in the California Sierra: implications for a warming climate. *J. Geophys. Res. Biogeosci.* **118**, 875–887 (2013).
 145. Hayashi, M. Alpine hydrogeology: The critical role of groundwater in sourcing the headwaters of the world. *Groundwater* **58**, 498–510 (2020).
 146. Earman, S., Campbell, A. R., Phillips, F. M. & Newman, B. D. Isotopic exchange between snow and atmospheric water vapor: estimation of the snowmelt component of groundwater recharge in the southwestern United States. *J. Geophys. Res.* **111**, D09302 (2006).
 147. Carroll, R. W. H., Deems, J. S., Niswonger, R., Schumer, R. & Williams, K. H. The importance of interflow to groundwater recharge in a snowmelt-dominated headwater basin. *Geophys. Res. Lett.* **46**, 5899–5908 (2019).
 148. Winnick, M. J. et al. Snowmelt controls on concentration-discharge relationships and the balance of oxidative and acid-base weathering fluxes in an alpine catchment, East River, Colorado. *Water Resour. Res.* **53**, 2507–2523 (2017).
 149. Dwire, K. A., Mellmann-Brown, S. & Gurrieri, J. T. Potential effects of climate change on riparian areas,

wetlands, and groundwater-dependent ecosystems in the Blue Mountains, Oregon, USA. *Clim. Serv.* **10**, 44–52 (2018).

150. Berghuijs, W. R., Woods, R. A. & Hrachowitz, M. A precipitation shift from snow to snowmelt leads to a decrease in streamflow. *Nat. Clim. Chang.* **4**, 583–586 (2014).

151. Davenport, F. V., Herrera-Estrada, J. E., Burke, M. & Duffenbaugh, N. S. Flood size increases nonlinearly across the western United States in response to lower snow-precipitation ratios. *Water Resour. Res.* **56**, e2019WR025571 (2020).

152. Kampf, S. K. & Lefsky, M. A. Transition of dominant peak flow source from snowmelt to rainfall along the Colorado Front Range: historical patterns, trends, and lessons from the 2013 Colorado Front Range floods. *Water Resour. Res.* **52**, 407–422 (2016).

153. Mallakpour, I., Sadegh, M. & AghaKouchak, A. A new normal for streamflow in California in a warming climate: wetter wet seasons and drier dry seasons. *J. Hydrol.* **567**, 203–211 (2018).

154. Barnhart, T. B. et al. Snowmelt rate dictates streamflow. *Geophys. Res. Lett.* **43**, 8006–8016 (2016).

155. Li, D., Lettenmaier, D. P., Margulis, S. A. & Andreadis, K. The role of rain-on-snow in flooding over the conterminous United States. *Water Resour. Res.* **55**, 8492–8513 (2019).

156. Hammond, J. C. & Kampf, S. K. Subannual streamflow responses to rainfall and snowmelt inputs in snow-dominated watersheds of the western United States. *Water Resour. Res.* **56**, e2019WR026132 (2020).

157. Godsey, S. E., Kirchner, J. W. & Tague, C. L. Effects of changes in winter snowpacks on summer low flows: case studies in the Sierra Nevada, California, USA. *Hydrol. Process.* **28**, 5048–5064 (2014).

158. Tague, C. & Grant, G. E. Groundwater dynamics mediate low-flow response to global warming in snow-dominated alpine regions. *Water Resour. Res.* **45**, W07421 (2009).

159. Safeeq, M., Grant, G. E., Lewis, S. L. & Tague, C. L. Coupling snowpack and groundwater dynamics to interpret historical streamflow trends in the western United States. *Hydrol. Process.* **27**, 655–668 (2013).

160. Trujillo, E., Molotch, N. P., Goulden, M. L., Kelly, A. E. & Bales, R. C. Elevation-dependent influence of snow accumulation on forest greening. *Nat. Geosci.* **5**, 705–709 (2012).

161. Hu, J., Moore, D. J. P., Burns, S. P. & Monson, R. K. Longer growing seasons lead to less carbon sequestration by a subalpine forest. *Glob. Change Biol.* **16**, 771–783 (2010).

162. Thorne, J. H. et al. The impact of climate change uncertainty on California's vegetation and adaptation management. *Ecosphere* **8**, e02021 (2017).

163. van Mantgem, P. J. et al. Widespread increase of tree mortality rates in the western United States. *Science* **323**, 521–524 (2009).

164. Allen, C. D. et al. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *For. Ecol. Manage.* **259**, 660–684 (2010).

165. Littell, J. S., McKenzie, D., Wan, H. Y. & Cushman, S. A. Climate change and future wildfire in the western United States: an ecological approach to nonstationarity. *Earths Future* **6**, 1097–1111 (2018).

166. Bentz, B. J. et al. Climate change and bark beetles of the western United States and Canada: direct and indirect effects. *Bioscience* **60**, 602–613 (2010).

167. Burns, S. P. et al. Snow temperature changes within a seasonal snowpack and their relationship to turbulent fluxes of sensible and latent heat. *J. Hydrometeorol.* **15**, 117–142 (2014).

168. Lundquist, J. D., Dickerson-Lange, S. E., Lutz, J. A. & Cristea, N. C. Lower forest density enhances snow retention in regions with warmer winters: a global framework developed from plot-scale observations and modeling. *Water Resour. Res.* **49**, 6356–6370 (2013).

169. Westerling, A. L., Hidalgo, H. G., Cayan, D. R. & Swetnam, T. W. Warming and earlier spring increase western US forest wildfire activity. *Science* **313**, 940–943 (2006).

170. Abatzoglou, J. T. & Williams, A. P. Impact of anthropogenic climate change on wildfire across western US forests. *Proc. Natl Acad. Sci. USA* **113**, 11770–11775 (2016).

171. Gleason, K. E., Nolin, A. W. & Roth, T. R. Charred forests increase snowmelt: effects of burned woody debris and incoming solar radiation on snow ablation. *Geophys. Res. Lett.* **40**, 4654–4661 (2013).

172. Staley, D. M. et al. Estimating post-fire debris-flow hazards prior to wildfire using a statistical analysis of historical distributions of fire severity from remote sensing data. *Int. J. Wildland Fire* **27**, 595–608 (2018).

173. Sengers, F. K. et al. Landslides after wildfire: initiation, magnitude, and mobility. *Landslides* **17**, 2631–2641 (2020).

174. Abraham, J., Dowling, K. & Florentine, S. Risk of post-fire metal mobilization into surface water resources: a review. *Sci. Total Environ.* **599–600**, 1740–1755 (2017).

175. Rust, A. J., Hogue, T. S., Saxe, S. & McCray, J. Post-fire water-quality response in the western United States. *Int. J. Wildland Fire* **27**, 203–216 (2018).

176. Wagenbrenner, J. W. & Robichaud, P. R. Post-fire bedload sediment delivery across spatial scales in the interior western United States. *Earth Surf. Process. Landf.* **39**, 865–876 (2014).

177. Olsen, W. H., Wagenbrenner, J. W. & Robichaud, P. R. Factors affecting connectivity and sediment yields following wildfire and post-fire salvage logging in California's Sierra Nevada. *Hydrol. Process.* **35**, e13984 (2021).

178. Smith, H. G., Sheridan, G. J., Lane, P. N. J., Nyman, P. & Haydon, S. Wildfire effects on water quality in forest catchments: a review with implications for water supply. *J. Hydrol.* **396**, 170–192 (2011).

179. Bladon, K. D., Emelko, M. B., Silins, U. & Stone, M. Wildfire and the future of water supply. *Environ. Sci. Technol.* **48**, 8936–8943 (2014).

180. Rhoades, C. C., Nunes, J. P., Silins, U. & Doerr, S. H. The influence of wildfire on water quality and watershed processes: new insights and remaining challenges. *Int. J. Wildland Fire* **28**, 721–725 (2019).

181. Maina, F. Z. & Siirila-Woodburn, E. R. Watersheds dynamics following wildfires: nonlinear feedbacks and implications on hydrologic responses. *Hydrol. Process.* **34**, 33–50 (2020).

182. Hallema, D. W. et al. Regional patterns of postwildfire streamflow response in the Western United States: the importance of scale-specific connectivity. *Hydrol. Process.* **31**, 2582–2598 (2017).

183. Hanan, E. J. et al. How climate change and fire exclusion drive wildfire regimes at actionable scales. *Environ. Res. Lett.* **16**, 024051 (2021).

184. Ho, M. et al. The future role of dams in the United States of America. *Water Resour. Res.* **53**, 982–998 (2017).

185. Luthy, R. G., Wolfand, J. M. & Bradshaw, J. L. Urban water revolution: sustainable water futures for California cities. *J. Environ. Eng.* **146**, 04020065 (2020).

186. Szinai, J. K., Deshmukh, R., Kammen, D. M. & Jones, A. D. Evaluating cross-sectoral impacts of climate change and adaptations on the energy-water nexus: a framework and California case study. *Environ. Res. Lett.* **15**, 124065 (2020).

187. Hartman, J., Steele, J., Frazier, S., Montgomery, A. *Recommendations for an Effective Water Rights Response to Climate Change* (State Water Resources Control Board, 2021).

188. Mateus, M. C. & Tullio, D. Reliability, sensitivity, and vulnerability of reservoir operations under climate change. *J. Water Resour. Plan. Manag.* **143**, 04016085 (2017).

189. The Metropolitan Water District of Southern California. Our foundation: securing our imported supplies. *mwdh20* <http://www.mwdh20.com/planning-for-tomorrow/securing-our-imported-supplies> (2021).

190. Elias, E. et al. Climate change, agriculture and water resources in the Southwestern United States. *J. Contemp. Water Res. Educ.* **158**, 46–61 (2016).

191. Hayhoe, K. et al. Emissions pathways, climate change, and impacts on California. *Proc. Natl Acad. Sci. USA* **101**, 12422–12427 (2004).

192. Balazs, C., Morello-Frosch, R., Hubbard, A. & Ray, I. Social disparities in nitrate-contaminated drinking water in California's San Joaquin Valley. *Environ. Health Perspect.* **119**, 1272–1278 (2011).

193. Gautam, M. R., Chief, K. & Smith, W. J. in *Climate Change and Indigenous Peoples in the United States: Impacts, Experiences and Actions* (eds Maldonado, J. K., Colombi, B. & Pandya, R.) 77–91 (Springer, 2014).

194. Doyle, J. T., Redsteer, M. H. & Eggers, M. J. Exploring effects of climate change on Northern Plains American Indian health. *Clim. Change* **120**, 643–655 (2013).

195. Madani, K., Guégan, M. & Uvo, C. B. Climate change impacts on high-elevation hydroelectricity in California. *J. Hydrol.* **510**, 153–163 (2014).

196. Vicuña, S., Dracup, J. A. & Dale, L. Climate change impacts on two high-elevation hydropower systems in California. *Clim. Change* **109**, 151–169 (2011).

197. Bartos, M. & Chester, M. Impacts of climate change on electric power supply in the Western United States. *Nat. Clim. Chang.* **5**, 748–752 (2015).

198. Averyt, K. et al. Sectoral contributions to surface water stress in the coterminous United States. *Environ. Res. Lett.* **8**, 035046 (2013).

199. Gleick, P. H. Water management: soft water paths. *Nature* **418**, 373 (2002).

200. Kaatz, L., Raucher, K. & Raucher, R. Embracing uncertainty: a case study examination of how climate change is shifting water utility planning (Stratus Consulting & Denver Water, 2015).

201. California Water Commission. Water storage investment program technical review: sites reservoir project. CWC <https://cwc.ca.gov/Water-Storage/WSIP-Project-Review-Portal/All-Projects/Sites-Project> (2018).

202. Hydro Review Content Directors. Reclamation accepting public comments on proposal to raise height of Shasta Dam. *Hydro Review* <https://www.hydroreview.com/dams-and-civil-structures/reclamation-accepting-public-comments-on-proposal-to-raise-height-of-shasta-dam/> (2020).

203. Perry, D. M. & Praskievicz, S. J. A new era of big infrastructure? (Re)developing water storage in the U.S. west in the context of climate change and environmental regulation. *Water Altern.* **10**, 437–454 (2017).

204. James, I. Utah's proposal to build Colorado River pipeline gets pushback from 6 states. *Arizona Republic* <https://www.azcentral.com/story/news/local/arizona-environment/2020/09/09/arizona-among-6-states-bucking-utahs-lake-powell-pipeline-proposal/5759035002/> (2020).

205. Ligon, F. K., Dietrich, W. E. & Trush, W. J. Downstream ecological effects of dams: a geomorphic perspective. *Bioscience* **45**, 183–192 (1995).

206. Liermann, C. R., Nilsson, C., Robertson, J. & Ng, R. Y. Implications of dam obstruction for global freshwater fish diversity. *Bioscience* **62**, 539–548 (2012).

207. Minear, J. T. & Kondolf, G. M. Estimating reservoir sedimentation rates at large spatial and temporal scales: a case study of California. *Water Resour. Res.* **45**, W12502 (2009).

208. Smith, R., Kasprzyk, J. R., Basdekas, L. & Dilling, L. Producing regionally-relevant multiobjective tradeoffs to engage with Colorado water managers [abstract H54F-08] (American Geophysical Union, 2016).

209. Smith, R., Kasprzyk, J. & Dilling, L. Participatory Framework for Assessment and Improvement of Tools (ParFAIT): increasing the impact and relevance of water management decision support research. *Environ. Model. Softw.* **95**, 432–446 (2017).

210. Sumargo, E., Cannon, F., Ralph, F. M. & Henn, B. Freezing level forecast error can consume reservoir flood control storage: potentials for Lake Oroville and New Bullards Bar reservoirs in California. *Water Resour. Res.* **56**, e2020WR027072 (2020).

211. Uccellini, L. W. & Ten Hoeve, J. E. Evolving the National Weather Service to build a weather-ready nation: connecting observations, forecasts, and warnings to decision-makers through impact-based decision support services. *Bull. Am. Meteorol. Soc.* **100**, 1923–1942 (2019).

212. California Department of Water Resources. 2020 Water resilience portfolio: in response to the Executive Order N-10-19 (California Department of Water Resources, 2020).

213. Luthy, R. G., Sharvelle, S. & Dillon, P. Urban stormwater to enhance water supply. *Environ. Sci. Technol.* **53**, 5534–5542 (2019).

214. Scanlon, B. R., Reedy, R. C., Faunt, C. C., Pool, D. & Uhlman, K. Enhancing drought resilience with conjunctive use and managed aquifer recharge in California and Arizona. *Environ. Res. Lett.* **11**, 035013 (2016).

215. Zhang, X. Conjunctive surface water and groundwater management under climate change. *Front. Environ. Sci.* **3**, 59 (2015).

216. Harpold, A. A. et al. Increasing the efficacy of forest thinning for snow using high-resolution modeling: a proof of concept in the Lake Tahoe Basin, California, USA. *Ecology* **13**, e2203 (2020).

217. Farooqi, T. J. A., Li, X., Yu, Z., Liu, S. & Sun, O. J. Reconciliation of research on forest carbon sequestration and water conservation. *J. For. Res.* **32**, 7–14 (2021).

218. Nelson, E. et al. Modeling multiple ecosystem services, biodiversity conservation, commodity production, and tradeoffs at landscape scales. *Front. Ecol. Environ.* **7**, 4–11 (2009).

219. French, J. R. et al. Precipitation formation from orographic cloud seeding. *Proc. Natl Acad. Sci. USA* **115**, 1168–1173 (2018).
220. Grant, S. B. et al. Taking the 'waste' out of 'wastewater' for human water security and ecosystem sustainability. *Science* **337**, 681–686 (2012).
221. Hering, J. G., Waite, T. D., Luthy, R. G., Drewes, J. E. & Sedlak, D. L. A changing framework for urban water systems. *Environ. Sci. Technol.* **47**, 10721–10726 (2013).
222. Cooley, H., Phurisamban, R. & Gleick, P. The cost of alternative urban water supply and efficiency options in California. *Environ. Res. Commun.* **1**, 042001 (2019).
223. Rao, P., Kostecki, R., Dale, L. & Gadgil, A. Technology and engineering of the water-energy nexus. *Annu. Rev. Environ. Resour.* **42**, 407–437 (2017).
224. Jägermeyr, J. et al. Integrated crop water management might sustainably halve the global food gap. *Environ. Res. Lett.* **11**, 025002 (2016).
225. Mitchell, J. et al. No-tillage and high-residue practices reduce soil water evaporation. *Calif. Agric.* **66**, 55–61 (2012).
226. Jägermeyr, J. et al. Water savings potentials of irrigation systems: global simulation of processes and linkages. *Hydrol. Earth Syst. Sci.* **19**, 3073–3091 (2015).
227. Davis, K. F., Seveso, A., Rulli, M. C. & D'Odorico, P. Water savings of crop redistribution in the United States. *Water* **9**, 83 (2017).
228. Gleick, P. H. et al. *Waste Not, Want Not: The Potential for Urban Water Conservation in California* (Pacifi Institute, 2003).
229. Schwabe, K., Nemati, M., Landry, C. & Zimmerman, G. Water markets in the Western United States: trends and opportunities. *Water* **12**, 233 (2020).
230. American Society of Civil Engineers. *The Economic Benefits of Investing in Water Infrastructure: How a Failure to Act Would Affect the US Economy Recovery* (American Society of Civil Engineers, 2020).
231. Harou, J. J. et al. Hydro-economic models: concepts, design, applications, and future prospects. *J. Hydrol.* **375**, 627–643 (2009).
232. Condon, L. E. & Maxwell, R. M. Groundwater-fed irrigation impacts spatially distributed temporal scaling behavior of the natural system: a spatio-temporal framework for understanding water management impacts. *Environ. Res. Lett.* **9**, 034009 (2014).
233. Lemos, M. C., Kirchhoff, C. J. & Ramprasad, V. Narrowing the climate information usability gap. *Nat. Clim. Chang.* **2**, 789–794 (2012).
234. McNie, E. C. Reconciling the supply of scientific information with user demands: an analysis of the problem and review of the literature. *Environ. Sci. Policy* **10**, 17–38 (2007).
235. Dewulf, A., Klenk, N., Wyborn, C. & Lemos, M. C. Usable environmental knowledge from the perspective of decision-making: the logics of consequentiality, appropriateness, and meaningfulness. *Curr. Opin. Environ. Sustain.* **42**, 1–6 (2020).
236. Flagg, J. A. & Kirchhoff, C. J. Context matters: context-related drivers of and barriers to climate information use. *Clim. Risk Manag.* **20**, 1–10 (2018).
237. Shi, L. et al. Roadmap towards justice in urban climate adaptation research. *Nat. Clim. Chang.* **6**, 131–137 (2016).
238. Araos, M. et al. Equity in adaptation: a systematic global review. *One Earth* https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3782859 (2021).
239. Anguelovski, I. et al. Equity impacts of urban land use planning for climate adaptation: critical perspectives from the global north and south. *J. Plan. Educ. Res.* **36**, 333–348 (2016).
240. Hawkins, E., Smith, R. S., Gregory, J. M. & Stainforth, D. A. Irreducible uncertainty in near-term climate projections. *Clim. Dyn.* **46**, 3807–3819 (2016).
241. Poff, N. L. et al. Sustainable water management under future uncertainty with eco-engineering decision scaling. *Nat. Clim. Chang.* **6**, 25–34 (2016).
242. Haasnoot, M., Kwakkel, J. H., Walker, W. E. & ter Maat, J. Dynamic adaptive policy pathways: a method for crafting robust decisions for a deeply uncertain world. *Glob. Environ. Change* **23**, 485–498 (2013).
243. Groves, D. G., Molina-Perez, E., Bloom, E. & Fischbach, J. R. in *Decision Making under Deep Uncertainty: From Theory to Practice* (eds Marchau, V. A. W. J., Walker, W. E., Bloemen, P. J. T. M. & Popper, S. W.) 135–163 (Springer, 2019).
244. Groves, D., Fischbach, J., Bloom, E., Knopman, D. & Keefe, R. *Adapting to a Changing Colorado River: Making Future Water Deliveries More Reliable Through Robust Management Strategies* (RAND Corporation, 2013).
245. Arnott, J. C., Mach, K. J. & Wong-Parodi, G. Editorial overview: The science of actionable knowledge. *Curr. Opin. Environ. Sustain.* **42**, A1–A5 (2020).
246. Meadow, A. M. et al. Moving toward the deliberate coproduction of climate science knowledge. *Weather Clim. Soc.* **7**, 179–191 (2015).
247. Cash, D. W. et al. Knowledge systems for sustainable development. *Proc. Natl Acad. Sci. USA* **100**, 8086–8091 (2003).
248. Dilling, L. & Lemos, M. C. Creating usable science: opportunities and constraints for climate knowledge use and their implications for science policy. *Glob. Environ. Change* **21**, 680–689 (2011).
249. Lemos, M. C. et al. To co-produce or not to co-produce. *Nat. Sustain.* **1**, 722–724 (2018).
250. Cash, D. et al. Saliency, credibility, legitimacy and boundaries: linking research, assessment and decision making. SSRN <https://doi.org/10.2139/ssrn.372280> (2002).
251. Cash, D. W., Borck, J. C. & Patt, A. G. Countering the loading-dock approach to linking science and decision making: comparative analysis of El Niño/Southern Oscillation (ENSO) forecasting systems. *Sci. Technol. Hum. Values* **31**, 465–494 (2006).
252. Goodrich, K. A. et al. Who are boundary spanners and how can we support them in making knowledge more actionable in sustainability fields? *Curr. Opin. Environ. Sustain.* **42**, 45–51 (2020).
253. Little, R. G. What's next for a national infrastructure policy: an encore or a requiem? *Public Work. Manag. Policy* **26**, 193–199 (2021).
254. Averyt, K. Are we successfully adapting science to climate change? *Bull. Am. Meteorol. Soc.* **91**, 723–726 (2010).
255. Zeng, X., Broxton, P. & Dawson, N. Snowpack change from 1982 to 2016 over conterminous United States. *Geophys. Res. Lett.* **45**, 12,940–12,947 (2018).
256. Gettelman, A., Morrison, H., Thayer-Calder, K. & Zarzycki, C. M. The impact of rimed ice hydrometeors on global and regional climate. *J. Adv. Model. Earth Syst.* **11**, 1543–1562 (2019).
257. Rhoades, A. M. et al. Sensitivity of mountain hydroclimate simulations in variable-resolution CESM to microphysics and horizontal resolution. *J. Adv. Model. Earth Syst.* **10**, 1357–1380 (2018).
258. van Kampenhout, L. et al. Improving the representation of polar snow and firn in the Community Earth System Model. *J. Adv. Model. Earth Syst.* **9**, 2583–2600 (2017).
259. Marsh, C. B., Pomeroy, J. W. & Spiteri, R. J. Implications of mountain shading on calculating energy for snowmelt using unstructured triangular meshes. *Hydrol. Process.* **26**, 1767–1778 (2012).
260. Lapo, K. E., Hinkelman, L. M., Raleigh, M. S. & Lundquist, J. D. Impact of errors in the downwelling irradiances on simulations of snow water equivalent, snow surface temperature, and the snow energy balance. *Water Resour. Res.* **51**, 1649–1670 (2015).
261. Fan, Y. et al. Hillslope hydrology in global change research and Earth system modeling. *Water Resour. Res.* **55**, 1737–1772 (2019).
262. Slater, A. G. et al. The representation of snow in land surface schemes: results from PILPS 2(d). *J. Hydrometeorol.* **2**, 7–25 (2001).
263. Jennings, K. S., Winchell, T. S., Livneh, B. & Molotch, N. P. Spatial variation of the rain–snow temperature threshold across the Northern Hemisphere. *Nat. Commun.* **11**, 48 (2019).
264. Fatchi, S. et al. Soil structure is an important omission in Earth System Models. *Nat. Commun.* **11**, 522 (2020).

Acknowledgements

P.S.N. and E.R.S.-W. acknowledge support from the Watershed Function Scientific Focus Area funded by the US Department of Energy, Office of Science, Office of Biological and Environmental Research under award no. DE-AC02-05CH11231. W.D.C. and A.M.R. are supported by the Director, Office of Science, Office of Biological and Environmental Research of the US Department of Energy through the Regional and Global Climate Modeling (RGCM) Program under the Calibrated and Systematic Characterization, Attribution and Detection of Extremes (CASCADE) Scientific Focus Area (award no. DE-AC02-05CH11231). A.D.J., A.M.R. and J.S. are supported by the Office of Science, Office of Biological and Environmental Research, Climate and Environmental Science Division of the US Department of Energy as part of the HyperFACETS Project. A Framework for Improving Analysis and Modeling of Earth System and Intersectoral Dynamics at Regional Scales (award no. DE-SC0016605). D.R.F. is supported by the Director, Office of Science, Office of Biological and Environmental Research of the US Department of Energy under contract no. DE-AC02-05CH11231 as part of their Atmospheric System Research (ASR) Program. C.T. is supported by the National Science Foundation's Hazard SEES program (award no. 1520847) and the National Science Foundation Critical Zone Observatory Network (award no. EAR-2012821). B.J.H. is supported by the Sulo and Aileen Maki Endowment. We thank M. Anderson, S. Hatchett, S. Hubbard, C. Koven and P. Ullrich for insightful conversations and constructive comments. We are grateful to W. Grimshaw, D. Yates and Denver Water for providing GIS data on conveyance, rivers and streamflows. We also thank D. Swantek for graphic design help.

Author contributions

E.R.S.-W. and A.M.R. initiated, led and contributed to all aspects of the Review. B.J.H. and L.S.H. contributed to the section on declining mountain snowpack. B.J.H., L.S.H. and W.D.C. contributed to the section on a low-to-no-snow future. C.T. contributed to the section on hydrological impacts. J.S., A.D.J., P.S.N. and L.K. contributed to the adaptation section. All authors contributed to the display items, the introduction and future directions sections, and participated in discussions, revisions and the final production of the manuscript.

Competing interests

The authors declare no competing interests.

Peer review information

Nature Reviews Earth & Environment thanks Ben Livneh, Philip Mote and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary information

The online version contains supplementary material available at <https://doi.org/10.1038/s43017-021-00219-y>.

© Springer Nature Limited 2021